

# *Journal of Mycology*

## TABLE OF CONTENTS

ATKINSON AND EDGERTON— <i>Protocoronospora</i> , a New Genus of Fungi.....	185
JENNINGS—A Case of Poisoning by <i>Amanita Phalloides</i> .....	187
DAVIS—A New Species of <i>Protomyces</i> .....	188
ARTHUR—Cultures of <i>Uredineae</i> in 1906.....	189
WILSON—An Historical Review of the Proposed Genera of <i>Phycomycetes</i> .....	205
KELLERMAN—Index to North American Mycology.....	210
KELLERMAN—Notes from Mycological Literature. XXV.....	228
EDITOR'S NOTES.....	232

**W. A. KELLERMAN, PH. D.**

*Professor of Botany, Ohio State University, Columbus, Ohio*

*Entered as Second Class Matter, Postoffice at Columbus, Ohio.*

## Cost of Separates.

---

Contributors who desire separates of their articles will receive the same at cost, approximately as follows:


For 4 pages or less:	100 copies	\$1 75	200 copies	\$2 25
" 8      "      "	100 "	2 50	200 "	3 25
" 16      "      "	100 "	4 25	450 "	5 50

For more extended articles proportionately higher.


Plates not included in the above.

**ADDRESS: EDITOR JOURNAL OF MYCOLOGY**





Digitized by the Internet Archive  
in 2025



*Elias J. Durand*

Journal of Mycology Portraits with Facsimile Autographs,



# Journal of Mycology

VOLUME 13—SEPTEMBER 1907

---

## TABLE OF CONTENTS

ATKINSON AND EDGERTON— <i>Protocoronospora</i> , a New Genus of Fungi.....	185
JENNINGS—A Case of Poisoning by <i>Amanita Phalloides</i> .....	187
DAVIS—A New Species of <i>Protomyces</i> .....	188
ARTHUR—Cultures of <i>Uredineae</i> in 1906.....	189
WILSON—An Historical Review of the Proposed Genera of <i>Phycomycetes</i> .....	205
KELLERMAN—Index to North American Mycology.....	210
KELLERMAN—Notes from Mycological Literature. XXV.....	228
EDITOR'S NOTES.....	232

---

## PROTOCOLONOSPORA, A NEW GENUS OF FUNGI.

(*Preliminary Note.*)

GEO. F. ATKINSON AND C. W. EDGERTON.

The senior author discovered a fungous disease of the cultivated vetch during July which does not seem to have been described before, and was first observed on the stems and pods from a small patch of vetch on the Horticultural grounds of Cornell University and later collected on vetch in the fields on the University farm where it seems to be abundant and a serious pest, often being associated with an *Ascochyta*. On the pods it often occurs quite pure, and here it is easily seen with the unaided eye to be distinct from the disease caused by the *Ascochyta*. It is, however, frequently mixed, even on the pods, with this fungus, but the very characteristic spots alone serve to distinguish it.

The spots are elongated, either narrow or elliptical, sometimes with a dull purple border. On the pods they are oblique. The spores ooze out in mass and have a pale pink or flesh color, but when spread in a thin layer, form a whitish film.

The fungus is subepidermal. The epidermis is ruptured in the form of a slit through which the spores escape. The mycelium becomes brown and then black, and the epidermis is later blackened; in age the spots are black oblique lines as seen on the pods, and many of them are sterile probably through failure of the fungus to fruit.

The structure of the fungus causing this new disease of vetch is very interesting. It resembles that of species of *Corticium*. The basidia form a definite hymenium which is seated on the pseudo-parenchymatous subhymenium, which is two or three cell-layers in thickness. The nourishing mycelium extends out into the surrounding tissue of the host. The spores are sessile; and are borne on a basidium in a whorl or crown at the end. The spores are oblong to subelliptical, straight or curved, continuous, hyaline, granular, and measure  $12-20 \times 3-3.5 \mu$ . As the spores fall away from the basidia others are produced as shown by cultures. Conidia similar to the basidiospores are produced on slender conidiophores which are intermingled with the basidia. This character recalls that of the genus *Exobasidium*. The spores also bud in yeast-like fashion from one or both ends, rarely from the side, and the sporidia thus produced are similar to the spores.

The fungus appears to be the type of a new genus for which the name *Protocoronospora* is proposed, and a provisional diagnosis is given as follows:

**Protocoronospora** Atkinson and Edgerton new genus. Stroma pseudoparenchymatous, two or three cell layers in thickness, formed by the compact branching of the mycelium, the ultimate exterior branches producing the basidia which form a hymenium. Spores sessile, hyaline, colorless, continuous, smooth, several (usually four-eight) on a basidium. Spores budding and forming sporidia similar in form. Conidia also similar in form on slender short conidiophores intermingled with the basidia.

**P. nigricans** Atkinson and Edgerton n. sp.—Forming narrow elongated spots on the pods, stems, leaves and bracts, spots oblique on the pods and from 2-5 mm. to 1-2 mm. Spots at first white or with purple border, later black. Stroma subepidermal, of pseudoparenchymatous cells  $6-9 \mu$  in diameter, two to three cell layers in thickness. Basidia clavate to subcylindrical,  $20-30 \times 6-8 \mu$ , 4-8 spored. Spores sessile, and basidia continuing to form new spores, at least in artificial culture. Spores pale pink in mass, oblong to subelliptical, hyaline, continuous, smooth, granular, straight or curved,  $12-20 \times 3-3.5 \mu$ , usually becoming once septate on germination. Mycelium from the stroma penetrating the adjacent tissues. Parasitic on pods, stems, leaves and bracts of *Vicia sativa*.

Botanical Department, Cornell University.  
September 2, 1907.



## A CASE OF POISONING BY AMANITA PHALLOIDES.

OTTO E. JENNINGS.

The writer's attention was recently called by Judge J. D. Shafer, of Pittsburg, to a newspaper account of a fatal case of mushroom poisoning at the little village of Deep Valley in the extreme southwestern point of Pennsylvania, and, acting upon Judge Shafer's urgent suggestion, the case was immediately investigated.

It was found that the village physician, Dr. Philip Dinsmore, together with three other members of the family and Mr. Frank Roberts, the man-of-all-work, had eaten with the evening meal, between six and seven o'clock, Sunday, August 4, a mess of mushrooms gathered that afternoon by Mr. Roberts. There had been about a quart of the mushrooms and they had been prepared by frying in flour and butter. All ate of the mushrooms excepting one little girl.

Between one and two o'clock the next morning all who had eaten of the mushrooms were taken violently sick, vomiting excessively and having an extreme diarrhoea. These symptoms continuing during Monday, Dr. H. C. Rice, of Freeport, Pa., was summoned and a treatment begun consisting of the sub-cutaneous injection of *atropine* and as far as possible the administration of narcotics and oleaginous purgatives.

The vomiting and diarrhoea continued for about three days, other symptoms being subnormal temperature, more or less delirium, and in the case of Dr. Dinsmore, severe muscular cramps of the limbs and extremities, and, evidently, of the muscular walls of the abdomen also, the patient dying early Thursday morning.

At the time of the writer's visit (Saturday, August 10) Mr. Roberts had so far recovered as to be about, but the other three patients were still confined to their beds. The vomiting and diarrhoea had ceased, but there was considerable enlargement of the liver with distension of the gall-bladder and the patients were becoming jaundiced.

Saturday morning Mr. Roberts escorted the writer to a little patch of about two acres of woods, lying at the base of the hillside along the creek, where the mushrooms had been gathered for the fatal meal. Two species were abundant, *Cantharellus*\* and the white form of *Amanita phalloides* Fr., and the latter species was indicated as the one composing the greater part of the mess taken. Other species indicated as having been also

---

Published by permission of Dr. W. J. Holland, Director of the Carnegie Museum.

\* The writer is indebted to Prof. D. R. Sumstine for verification of the identifications.

selected were *Amanitopsis vaginata* (Bull.) Roz., and *Russula emetica* Fr. — a very few. The only test applied in selecting the fungi had apparently been the pleasing appearance and the tenderness of the mushroom. Roberts' identification of *Amanita* as composing the greater part of those eaten was independently verified by one of the patients, Dr. Dinsmore's sister, who had prepared the fungi for eating.

From the evidence obtained it is quite clear that the poisoning was due to the deadly *Amanita*, and it will be noticed that the symptoms exhibited were in close agreement with those ascribed to *phallin* poisoning by chestnut,\*\* although Dr. Rice characterized the intestinal discharges as "serous" and not assuming the "rice-water" condition, and neither extreme salivation nor decided suppression of the urine was noticed.

In connection with the supposed action of *phallin* in decomposing the blood corpuscles and in bringing about the escape of the blood serum from the system by way of the alimentary canal it may be mentioned as a partial confirmation that the undertaker experienced considerable trouble in preparing the corpse for burial,—less than half the usual amount of blood could be extracted; thus indicating a depletion of blood supply before death occurred.

CARNEGIE MUSEUM, August 14, 1907.

(Published by permission of Dr. W. J. Holland, Director.)

---

\*\* Chestnut, V. K. Circular No. 13, Div. Botany, U. S. Dept. Agriculture.

---

## A NEW SPECIES OF PROTOMYCES.

J. J. DAVIS.

For the purpose of securing a name under which to distribute specimens in *Fungi Columbiani* I submit the following:

**Protomyces gravidus n. sp.**—Causing hypertrophic swellings on stems, branches, petioles and midribs. Spores, either sub-epidermal or in the vascular bundles but not usually in both, numerous, surface more or less irregularly uneven, generally globose but some times elliptical, ovate or polygonal, 30-55 x 27-40  $\mu$ , plurinucleate; epispore thin (1-3  $\mu$ ), brown; endospore in maturity thick (3-5 $\mu$ ), hyaline. On *Bidens cernua* L. and *Bidens connata* Muhl., Dousman; on the same hosts and sparingly on *Bidens frondosa* L., Racine; on *Bidens cernua* L., Berryville, all in Wisconsin. July to November.

What I have called the endospore should rather perhaps be considered a peripheral layer of cytoplasm in a resting condition the true endospore being a hyaline membrane 1  $\mu$  or less thick.



In the 35th Rep. of the New York State Botanist, p. 138, is reported the occurrence of *Protomyces macrosporus* Ung. on leaves and stems of *Ambrosia trifida* L., at Albany, with a brief description that corresponds with the fungus on *Bidens*. Prof. Peck informs me that it was abundant at one station during one season. Through the kindness of Dr. Farlow I have had an opportunity to examine sections of a gall on *Ambrosia artemisiaefolia* L. which was sent him from Nantucket, Mass. in August 1905, containing spores similar to those in the *Bidens* galls.

Sydow described in *Annales Mycologici*, 1:237, *Entyloma leucanthemi* which was distributed by Vestergrén (*Microm.* No. 808.) under the name *Protomycopsis leucanthemi* (Syd.) Magn. but I have been unable to learn of a publication of the characters of the genus. Again through the kindness of Dr. Farlow I have been able to examine sections containing this fungus. The spores are similar to those in *Bidens* but they appear to occur in the leaf blade without gall formation and no mention is made of such swellings by Sydow. I therefore hesitate to distribute my material under the name given by Magnus.

I have found the fungus here considered on no hosts other than *Bidens* — not even on *Coreopsis* growing with affected *Bidens* — and for my present purpose the question as to the relation between the *Ambrosia* and *Bidens* inhabiting forms may be left open.

I have made many attempts to observe the germination of the spores, at all seasons of the year, using material kept continuously out of doors but without result.

Racine, Wisconsin, August 6, 1907.

---

## CULTURES OF UREDINEAE IN 1906.<sup>1</sup>

BY J. C. ARTHUR.

The present article forms the seventh of a series of reports<sup>2</sup> by the author upon the culture of plant rusts, covering the years from 1899 to the close of 1906. As in previous years the grass and sedge rusts have constituted a large part of the list of species under trial. This is partly due to the economic and scientific interest connected with them, but even more, possibly, to the greater ease with which wintered-over and viable spores may be secured for cultural study. Among the species whose life-

<sup>1</sup> Read before the Botanical Society of America at the New York meeting, December 31, 1906.

<sup>2</sup> See Bot. Gaz. 29:268-276, 35:10-23; Jour. Myc. 8:51-56, 10:8-21, 11:50-67 and 12:11-27.

cycle has now been worked out for the first time, the one having the greatest economic import does not belong to the grass or sedge forms, but inhabits flax. The discovery of the full developmental history of this serious menace to successful flax growing, brought about by the cooperation of Professor Henry L. Bolley of North Dakota, widely known for his interest and scientific studies in promoting the flax industry, has been a matter of much satisfaction. It greatly clarifies the problem of controlling the flax rust in the interest of the cultivator.

The work of testing the viability of spores, making the sowings, and recorded data, required, as in previous years, the undivided attention of a person to whom the whole work could be intrusted. The expense of such an assistant was this year met by a grant from the Botanical Society of America. It was the second time the Society has given material aid to this series of investigations.

Through the kindness of Professor R. A. Harper, I was so fortunate as to enlist the interest of Dr. E. W. Olive, lecturer at the University of Wisconsin, who consented to supervise the season's work. The forethought and constant watchfulness, the enthusiastic application, and especially the maturity of judgment and breadth of knowledge brought to bear on the work by Dr. Olive materially increased the completeness of the results.

As in former years correspondents have provided much of the material used in the trials, partly upon their own initiative, and partly in response to suggestion, for all of which I am under heavy obligation. Teliosporic material was sent by Messrs. A. D. Selby, Wooster, Ohio; H. H. Whetzel, Ithaca, N. Y.; Chas. E. Fairman, Lyndonville, N. Y.; W. A. Kellerman, Columbus, Ohio; H. L. Shantz, Columbia, Mo.; J. J. Davis, Racine, Wis.; John L. Sheldon, Morgantown, W. Va.; H. L. Bolley, Agricultural College, N. D.; E. Bartholomew, Stockton, Kans.; Geo. E. Morris, Waltham, Mass.; Guy W. Wilson, Lafayette, Ind.; E. Bethel, Denver, Colo.; and especially by Rev. J. M. Bates, Red Cloud, Neb. Aeciosporic material was sent by Messrs. Herman von Schrenk, St. Louis, Mo.; H. H. Whetzel, Ithaca, N. Y.; C. L. Shear, Tacoma Park, D. C.; D. Reddick, Ithaca, N. Y.; and P. H. Rolfs, Lake City, Fla., all the collections being either *Caeoma* or *Peridermium* on species of *Pinus*. Host plants are often required for the work, which do not grow in this vicinity, and can not be purchased from dealers, and for a number of such plants in good growing condition I am indebted to Messrs. William Trelease of the Missouri Botanical Garden, St. Louis, Mo.; John L. Sheldon, Morgantown, W. Va.; P. B. Kennedy, Reno, Nev.; and J. J. Davis, Racine, Wis.

During the present season 94 collections of material with resting spores and 15 collections with active spores were employed, from which 293 drop cultures and 6 Petri dish cultures



were made to test the germinating condition of the spores. Out of the 94 collections with resting spores 46 could not be made to germinate, although no reason could be assigned why they should not. This gave 48 collections of available material belonging to 30 species of rusts, exclusive of the aecial pine rusts, and from these 223 sowings were made. Beside these 53 sowings were made with *Caeoma* and *Peridermium* spores from pine, all without infection, 27 sowings with teliospores of *Gymnosporangium*, and 23 sowings with various aeciospores. Altogether 326 sowings were made, and for this purpose 134 species of hosts were required, which were grown temporarily in the greenhouse, where practically all the work was done. The results of this work are given in the following paragraphs, and are divided into negative results, positive results with species whose life histories have already been previously determined, and positive results with species whose life histories have not before been fully known.

---

Of the trials giving negative results the following may be recorded to serve for reference in further studies.

1. PUCCINIA on *Carex Pennsylvanica* Lam., collected near Lafayette, Ind., was sown on *Trillium recurvatum*, *Napaea dioica*, *Anemone thalictroides*, *Isopyrum bitermum*, *Anemone virginiana*, *Actaea alba*, *Viola cucullata*, *Dirca palustris*, *Polemonium reptans*, *Ambrosia trifida*, *Rudbeckia laciniata*, and *Lactuca canadensis*, with no infection. Similar material in former seasons has been tried on eighteen other species of hosts with negative results.<sup>3</sup>

2. PUCCINIA on *Carex gravior* Bailey, sent by Rev. J. M. Bates from Red Cloud, Neb., was sown on *Actaea rubra*, *Thalictrum dioicum*, *Isopyrum bitermum*, *Apios Apios*, *Falcata comosa*, *Psoralea Onobrychis*, *Cassia Chamaecrista*, *Polygala Senega*, *Aesculus glabra*, *Ceanothus americanus*, *Smilax herbacea*, *Viola cucullata*, *Napaea dioica*, *Callirrhoe involucrata*, *Althaea rosea*, *Hibiscus Moscheutos*, *Macrocalyx Nyctelea*, *Polemonium reptans*, *Myosotis palustris*, *Phlox divaricata*, *Phlox subulata*, *Triosteum perfoliatum*, *Boltonia asteroides*, *Laciniaria pycnostachya*, *Rudbeckia laciniata*, *Senecio obovatus*, and *Cacalia reniformis*, with no infection. Similar material from the same source has been sown in previous years upon eleven other species of hosts with negative results.<sup>4</sup>

3. PUCCINIA on *Polygonum scandens* L., obtained in the vicinity of Lafayette, Ind., was sown five times on *Geranium maculatum*, twice on *G. Robertianum*, twice on *G. pusillum*, and

---

<sup>3</sup> See Jour. Myc. 10:10. 1904; 11:51. 1905; and 12:12. 1906.

<sup>4</sup> See Jour. Myc. 10:10. 1904; and 11:52. 1905.

twice on *Thalictrum dioicum*. These sowings were made under seemingly favorable conditions and yet no infection resulted. In 1903 Dr. W. Tranzschel of St. Petersburg established the connection between *Puccinia Polygoni-amphibii* Pers. on *Polygonum amphibium* and the aecia on *Geranium palustre* and *G. pratense*, and a year later the writer corroborated the discovery with corresponding American species of hosts. In 1904 Dr. Tranzschel showed that the rust on climbing species of *Polygonum*, often included with the preceding, is distinct, either as a true species or a biological species, for it produces its aecia on *Geranium pusillum*. To see if this also could be substantiated with American material the above sowings were made with seemingly excellent teliosporic material, but the negative results leave the matter an open question. The only other native *Geranium* on which this form might be expected to grow readily is *G. carolinianum*, which was unfortunately not at hand for the test.

4. PUCCINIA ON *Muhlenbergia diffusa* Schreb., sent by Rev. J. M. Bates from Red Cloud, Neb., was sown on *Trillium recurvatum*, *Actaea alba*, *Anemonella thalictroides*, *Isopyrum biter-natum*, *Caulophyllum thalictroides*, *Apios Apios*, *Viola pubescens*, *Dirca palustris*, *Althaea rosea*, *Callirrhoe involucrata*, *Napaea dioica*, *Hibiscus Moscheutos*, *Marcrocalyx Nyctelea*, *Polemonium reptans*, *Ambrosia trifida*, and *Lactuca canadensis*, with no infection. This taken with previous trials shows that the rusts on different species of *Muhlenbergia* are in all probability biologically complex.

5. PUCCINIA SCHEDONNARDI K. & S., sent by Rev. J. M. Bates from Red Cloud, Neb., was sown on *Callirrhoe involucrata*, *Althaea rosea*, and *Ceanothus americanus*, with no infection. Like material from the same source was sown in 1902 on eight other species of hosts with negative results.<sup>5</sup> The small sori and fine leaves of the grass make the manipulation of material of this species somewhat uncertain.

6. PUCCINIA EMACULATA Schw. on *Panicum capillare*, obtained in the vicinity of Lafayette, Ind., where it is very common, was sown on *Polygala Senega* and *Napaea dioica*. This rust was sown in previous seasons on eighteen other species of hosts.<sup>6</sup>

7. UROMYCES on *Juncus effusus* L., sent by Dr. Charles E. Fairman from Ridgeway, N. Y., was sown on *Polemonium reptans*, *Houstonia purpurea*, *Ambrosia trifida*, *Rudbeckia laciniata*, *Polymnia canadensis*, *Parthenium integrifolium*, *Silphium integrifolium*, *S. perfoliatum* *S. terebinthinaceum*, and *Senecio obovatus*,

<sup>5</sup> See Bot. Gaz. 35:11. 1903.

<sup>6</sup> See Bot. Gaz. 35:12. 1903; Jour. Myc. 8:52. 1902; 10:10. 1904; and 12:12. 1906.



with no infection. What was doubtless the same rust, and also from western New York, was sown in 1905 on two other species of hosts with negative results.<sup>7</sup>

This rust has heretofore been considered to belong to *Uromyces Junci* (Desm.) Tul., but recent study has shown that it is morphologically quite distinct from that species, especially as it has urediniospores that are echinulate and four-pored, instead of verrucose and two-pored, as in the European species, which by the way apparently does not occur in the United States east of Nebraska and Kansas. It was described by Schweinitz (Trans. Am. Phil. Soc. 4:295. 1832.) as a new species under the name *Puccinia Junci*. As that specific name is not now available, I suggest that the species be called *Uromyces effusus*, in allusion to the copious distribution of the sori over the surface of the host, and would characterize it as follows:

***Uromyces effusus* sp. nov.**

O and I. Pycnia and aecia unknown.

II. Uredinia amphigenous, scattered, oblong or linear, 0.1-0.3 mm. wide by 0.3-1.5 mm. long, tardily naked, dark cinnamon-brown, ruptured epidermis very conspicuous; urediniospores broadly ellipsoid or oval, 14-19 by 18-26  $\mu$ , wall light yellow about 1.5  $\mu$  thick, rather sparingly and bluntly echinulate, pores 4, equatorial.

III. Telia amphigenous, numerous, scattered, oblong or linear, 0.2-0.5 mm. wide by 0.3-2 mm. or more long, rarely confluent, finally naked, ruptured epidermis very conspicuous; teliospores obovate or broadly oval, 13-19 by 24-33  $\mu$ , obtuse or rarely acute at apex, usually narrowed below; wall chestnut-brown, 1.5-2  $\mu$  thick, much thicker above, 6-10  $\mu$ , smooth; pedicel tinted, about as long as the spore.

On *Juncus effusus* L. Type collected by L. von Schweinitz at Bethlehem, Pa. Collections in the writer's herbarium from Ohio, New York, West Virginia, and Maryland, and in the herbarium of the New York Botanical Garden from New Jersey, Massachusetts and South Carolina. It is also found in the following exsiccati: Ellis, N. Am. Fungi. 238; Ellis & Ev., Fungi Columb. 339; Ravenel, Fungi Am. 51; Shear, N. Y. Fungi 76; Kellerm., Ohio Fungi 38.

8. UROMYCES ELEOCHARIDIS Arth. on *Eleocharis palustris* (L.) R. & S., sent by Mr. E. Bartholomew from Stockton, Kan., was sown on *Callirrhoe involucrata*, *Napaea dioica*, *Cassia chamaecrista*, *Myosotis palustris*, and *Silphium perfoliatum*, with no infection.

9. UROMYCES ACUMINATUS Arth. on *Spartina cynosuroides* Willd., obtained at Fair Oaks, Ind. by Mr. F. D. Kern, was sown four times on *Steironema ciliatum*, twice on *S. lanceolatum*, twice on *Lysimachia quadrifolia* L., and once each on *L. terrestris*, *Polygala Senega*, *Napaea dioica* and *Houstonia purpurea*, with no infection. As teliosporic material of this rust, obtained from Nebraska, was sown on *Steironema ciliatum* with success in 1905,<sup>8</sup>

<sup>7</sup> See Jour. Myc. 12:13. 1906.

<sup>8</sup> See Jour. Myc. 12:24. 1906.

it would seem that what now passes under one name is either a segregate, or a series of biological species, and that the *Uromyces* on *Spartina* in Indiana differs in some way from that in Nebraska.

---

The following species of rusts were successfully grown, and the data supplement that obtained from previous cultures of this series, or that recorded by other American or European investigators.

1. *MELAMPORA BIGELOWII* Thuem. — Teliosporic material obtained near Lafayette, Ind., on *Salix* sp., was sown April 25 on *Larix decidua*, pycnia appearing in abundance May 2, and fully grown aecia about May 12.<sup>9</sup>

2. *CRONARTIUM QUERCUS* (Brond.) Schroet. — Work with this species was suggested by Dr. C. L. Shear, who also provided freshly gathered aecia on *Pinus virginiana* Mill., sending these a number of times in varying quantity. Dr. Shear made a number of cultures in the open in the spring of 1905, and presented a paper embodying his observations and conclusions<sup>10</sup> at the New Orleans meeting of the American Mycological Society, but which did not appear in print until June, 1906, after all data to be presented here were secured.

Aeciospores from material provided by Dr. Shear was sown May 12 in the greenhouse on three plants of *Quercus alba* and two plants of *Q. velutina*. On May 19 another sowing was made on two other plants of *Q. velutina*. This work coming late in the season did not receive daily examination, but on June 1 all the plants of *Q. velutina* showed uredinia, and one of them also had developed telia. By June 25 the remaining four plants had produced telia. The plants of *Q. alba* remained free from infection.

The aecia used for these cultures were the typical form of *Peridermium Cerebrum* Peck. There seems no reason to doubt the identity of the American, European and Japanese fungus, which has passed under a number of names, but a discussion of the literature and facts will not be taken up in this place.

3. *PUCCINIA OPIZII* Bubák. — Aecia on various wild species of *Lactuca*, and even on the garden *L. sativa*, are common in the extended region of the upper Mississippi valley. They were described by Burrill (Bull. Ill. Lab. Nat. Hist. 2: 232. 1885), but the name *Aecidium compositarum Lactucæ* Burr. was first published three years later (Saccardo, Syll. fung. 7: 799. 1888). This form has often appeared in considerable abundance within

<sup>9</sup> For previous cultures see Jour. Myc. 11:60. 1905.

<sup>10</sup> Jour. Myc. 12:89-92. 1906.



a hundred feet of my laboratory door, and much attention has been bestowed upon it. The first clue to its connection was found on the last day of April of this year, when in company with Mr. F. D. Kern, the writer detected at Fair Oaks, Ind., a hundred miles north of this place, in two well separated spots, some plants of *Lactuca canadensis* thickly covered with aecia, and intermixed with the affected leaves some leaves of a small, narrow leaved *Carex*, bearing teliospores of the previous year's growth. No evidence of fruiting could be found on the *Carex*, and roots brought back and grown in pots have shown no signs of fruit, so that the *Carex* has not been specifically determined. Teliosporic material was obtained from both localities, and May 2 one was sown on *Lactuca canadensis* and *Onagra biennis*, the other on *L. virosa* and two plants of *L. canadensis*. *Onagra* showed no infection, but all plants of *Lactuca* gave rise to pycnia May 9, and aecia May 15, in great abundance. Another sowing was made May 14 on *L. sativa*, which gave pycnia May 14, and aecia May 28, with ample development.

This rust is taken to be the same as the one which Dr. Fr. Bubák studied in Bohemia by means of cultures, and which he has very fully described.<sup>11</sup> No Bohemian collections, however, are at hand with which to make comparison, but two European collections of aecia (Sydow, Uredineen 334 and 1100) show essential morphological agreement with American aecia on *Lactuca*. Dr. Bubák (l. c.) has stated that to him the American and European collections appear distinct, but without saying wherein the difference may lie. To me the differences appear to be habitual. On thin leaved hosts both pycnia and aecia are in more open and indefinite groups. On hosts from the western prairies, which have firm and strongly developed leaves the groups of aecia are usually compact and circumscribed, and surround the often amphigenous pycnia. The European aecia belonging to the species are known under the name *Accidium lactucinum* Lagerh. & Lindr.

4. PUCCINIA SAMBUCI (Schw.) Arth. — Teliosporic material on *Carex Frankii* Kunth, brought from Frankfort, Ind., by Mr. F. D. Kern, was sown May 10 on *Sambucus canadensis*, giving numerous pycnia May 16, and abundance of aecia May 26. This adds another host to this common species, those already known being *Carex trichocarpa*, *C. lurida*, and *C. lupulina*.<sup>12</sup>

5. PUCCINIA PECKII (DeT.) Kellerm. — Teliosporic material on *Carex trichocarpa* Muhl., brought from Fair Oaks, Ind., was sown on *Onagra biennis* May 4, and gave rise to pycnia May 14, and to aecia May 17, both in abundance. Another collection on *C. lanuginosa* Michx., sent by Rev. J. M. Bates

<sup>11</sup> Centr. Bakt. 9<sup>2</sup>:924. 1902.

<sup>12</sup> See Bot. Gaz. 35:14. 1903; Jour. Myc. 8:55. 1902; 12:14. 1906.

from Wymore, Neb., was sown on *Onagra biennis* May 19, giving rise to pycnia May 26, and aecia June 2, both in abundance.<sup>13</sup>

6. *PUCCINIA ALBIPERIDIA* Arth. — This rust on three species of hosts was obtained in different localities near Lafayette, Ind., and sown with the following results:

From *Carex squarrosa* L., sown in greenhouse April 16 on *Ribes rotundifolium*; April 21, pycnia; April 30, aecia.

From *C. squarrosa* L., sown in garden April 21 on *R. gracile*; April 25, pycnia; May 13, aecia.

From *C. squarrosa* L., sown in greenhouse April 16 on *R. rubrum*; no infection.

From *C. tetanica* Schk., sown in greenhouse April 20 on *R. Cynosbati*; April 27, pycnia; May 9, aecia.

From *C. crinita* Lam., sown on *R. Cynosbati* in greenhouse April 26, then plant transferred to garden; May 4, pycnia; May 17, aecia.

These results add one more telial host to those previously used for cultures.<sup>14</sup> They also have given an opportunity for a study of the differences between the pale aecia obtained by cultures and the highly colored aecia usually observed in the field. The aecia grown wholly in the greenhouse were pale, as in previous years; those on the plant which had the pot plunged into the garden soil after the fungus became established, were much more colored; and those raised from sowings made in the garden were highly colored and presented essentially the same appearance as others that came upon some nearby bushes of *Ribes* from natural infection. The result of observations during the last six years, coupled with the cultures of this year, make the conclusion almost inevitable that shade, moist air, and slow growth, tend to make the aecia smaller, with less coloring matter in the peridial cells and surrounding mycelium, and also tend to produce less hypertrophy of the tissues of the host, and that this accounts for the differences observed between aecia grown in cultures and those very common on *Ribes Cynosbati*, *R. rotundifolium*, *R. gracile*, and similar species of gooseberries throughout the eastern United States. All collections of this sort, therefore, may be called *Puccinia albiperidia*, but whether this is a distinct species from the very similar rust of Europe, *Puccinia Grossulariae* (Schum.) Lagerh., or one of the several biological species established by Klebahn, still remains an open question.

7. *PUCCINIA ANGUSTATA* Peck. — Teliosporic material on *Scirpus atrovirens* Muhl., from the vicinity of Lafayette, Ind., was sown April 28, on *Dirca palustris*, with no infection. On

<sup>13</sup> For previous cultures see Bot. Gaz. 35:13. 1903; Jour. Myc. 8:55. 1902; 11:58. 1905; and 12:15. 1906.

<sup>14</sup> For previous cultures see Jour. Myc. 8:53. 1902; 10:11. 1904; 11:58. 1905; and 12:14. 1906.



May 4 it was sown on *Lycopus Americanus*, giving rise May 14 to numerous pycnia, and May 18 to aecia in abundance.<sup>15</sup>

8. PUCCINIA ELEOCHARIDIS Arth. — Teliosporic material on *Eleocharis palustris* (L.) R. & S., sent by Mr. E. Bartholomew from Stockton, Kans., was sown May 7 on *Eupatorium perfoliatum*, giving rise to pycnia May 14, and to aecia May 22. Similar material on same species of host from near Lafayette, Ind., was sown June 1 on *E. perfoliatum*, giving rise to pycnia June 8, and to aecia June 20. It was also sown on *Napaea dioica*, with no infection.<sup>16</sup>

9. PUCCINIA ANDROPOGONIS Schw. — Teliosporic material on *Andropogon scoparius* Michx., sent by Rev. J. M. Bates from Sargent, Neb., was sown April 24 on *Pentstemon hirsutus*, giving rise to pycnia April 30, and to aecia May 10, both in abundance.<sup>17</sup>

10. PUCCINIA TOMIPARA Trel. — Teliosporic material on *Bromus purgans* L., from Lafayette, Ind., was sown May 19 on *Clematis virginiana*, giving rise to pycnia May 26, and to aecia June 8, both in abundance.<sup>18</sup>

11. PUCCINIA SUBNITENS Diet. — Teliosporic material on *Distichlis spicata* (L.) Greene, sent by Rev. J. M. Bates from Red Cloud, Neb., was sown April 5 on *Chenopodium album*, *Bursa Bursa-pastoris*, and *Sarcobatus vermiculatus*, giving a weak infection only on the *Chenopodium*. As the *Sarcobatus* plant soon died, another sowing was made May 5 on two other plants of *Sarcobatus*, but with no infection. It was sown again May 10, and May 19, on *Sarcobatus*, still with no infection. Still a fifth sowing was made May 29 on two plants of *Sarcobatus*, and one of *Chenopodium album*, with a weak infection of the latter, and with apparently a few pycnia showing on one leaf of *Sarcobatus*. The plants of *Sarcobatus* were sent by Dr. P. B. Kennedy from Reno, Nev., and had not become established when the sowings were made. They were obtained where *Distichlis spicata* grew intermixed, well covered with *Puccinia subnitens*, and the *Sarcobatus* was well besprinkled with aecia, not distinguishable from those now known to belong to this grass rust.<sup>19</sup> The particular object in view was to determine experimentally if *Puccinia subnitens* will grow on *Sarcobatus*. The single seeming infection is doubtful, as it may have come from spores transported with the plants. The question remains an open one,

<sup>15</sup> For previous cultures see Bot. Gaz. 29:273. 1900; and Jour. Myc. 8:53. 1902.

<sup>16</sup> For previous cultures see Jour. Myc. 12:23. 1906.

<sup>17</sup> For previous cultures see Bot. Gaz. 29:272. 1900; Jour. Myc. 9:10. 1903; and 10:11. 1904.

<sup>18</sup> For previous cultures see Jour. Myc. 11:62. 1905.

<sup>19</sup> For previous cultures see Bot. Gaz. 35: 19. 1903; Jour. Myc. 11:54. 1905; 12:16. 1906.

although I venture the opinion that if the teliosporic material had come from Nevada, instead of Nebraska, the sowings would have been successful.

12. *PUCCINIA POCULIFORMIS* (Jacq.) Wettst. — Teliosporic material on *Agrostis alba* L., brought from Fair Oaks, Ind., was sown May 2 on two plants of *Berberis vulgaris*, both showing abundant pycnia May 9, and aecia May 18.

Aeciospores from these cultures were sown May 31 on *Avena sativa*, *Hordeum vulgare* (Great Beardless), and *Triticum vulgare* (Jones' Silver Sheaf), without infection in the first case, and with sparing infection in the other two cases, the former showing uredinia June 12, and the latter somewhat later.

Teliosporic material on *Elymus canadensis* L., sent from Racine, Wis., by Dr. J. J. Davis, was sown May 2 on *Berberis vulgaris*, showing pycnia May 9, and aecia May 18. Aecia from this culture were sown May 31 on *Triticum vulgare* (Jones' Silver Sheaf), and *Secale cereale* (Mammoth Winter Rye), with no infection.<sup>20</sup>

13. *PUCCINIA TRANSFORMANS* Ellis & Ev. — Teliosporic material from a greenhouse plant of *Stenolobium Stans*, which had been infected a year previously, was sown May 7 on two healthy plants of the same species, and gave rise to pycnia May 22, and to telia May 30. A sowing on two other plants was made May 10, showing pycnia May 26, and telia May 31.<sup>21</sup>

14. *PUCCINIA XANTHII* Schw. — Teliosporic material on *Xanthium* sp., obtained in the vicinity of Lafayette, Ind., about the middle of April, was sown on *Xanthium* seedlings April 20, and gave rise to translucent papillae simulating pycnia April 26, and open telia May 3. Another sowing May 1 gave pale papillae about May 18, and open telia about May 26.<sup>22</sup>

15. *PUCCINIA SILPHII* Schw. — Teliosporic material on *Silphium integrifolium* Michx., obtained the last of March near Lafayette, Ind., was sown April 26 on *Silphium perfoliatum*, giving rise to pale papillae May 1, and open telia May 4. Another sowing was made on *S. terebinthinaceum* May 1, giving rise to pale papillae May 8, and open telia May 14. Similar material from another locality was sown May 3 on *S. integrifolium*, *S. perfoliatum* and *S. terebinthinaceum*, giving infection in usual way in each instance but exact data not taken.<sup>23</sup> It was observed that the infection on *S. integrifolium* was more rapid in its growth and more abundant than on the other hosts. In the report of last

<sup>20</sup> For previous cultures see Jour. Myc. 8:53. 1902; 11:57. 1905; 12:17. 1906.

<sup>21</sup> For previous cultures see Jour. Myc. 12:22. 1906.

<sup>22</sup> For previous cultures see Jour. Myc. 12:20. 1906.

<sup>23</sup> For previous cultures see Jour. Myc. 12:21. 1906.



year's cultures the opinion was expressed that this rust may be composed of biological races, but the present work shows that adaptation to the hosts is not so close but that under specially favorable conditions the rust may be transferred from one host to another.

16. PUCCINIA PRUNI-SPINOSAE Pers.—Aecia on *Hepatica acutiloba* DC., from near Lafayette, Ind., were sown April 28 on *Prunus serotina* and *Amygdalus Persica*, giving rise May 21 to uredinia on the former host, but with no infection on the latter host. Like material was sown May 2 on *Prunus serotina* and *P. pumila*, giving rise to the uredinia in both instances May 21. These results abundantly confirm, and somewhat extend, the work of last year.<sup>24</sup>

17. UROMYCES SCIRPI (Cast.) Burr.—Teliosporic material on *Scirpus fluviatilis* (Torr.) A. Gray, sent by Rev. J. M. Bates from Walbach, Neb., was sown May 22 on *Cicuta maculata*, giving abundance of pycnia May 31, and of aecia June 8. A sowing made June 1 on *Pastinaca sativa*, the plants being especially thrifty, gave no infection. On June 5 another sowing was made on *Oxypolis rigidus* and *Cicuta maculata*, with no infection on the former, but with fine showing of pycnia on the latter June 12, and of aecia June 22.

There is apparently no morphological difference between the American rust and the corresponding European one, and the hosts are also much alike. Sixteen years ago Dr. P. Dietel showed by cultures<sup>25</sup> that in central Germany aecia are produced on *Sium latifolium*, which in habit and structure is much like *Cicuta maculata*. He also found that, most curiously, aecia could be grown from the same material on *Hippurus vulgaris*, which belongs to another family of plants, showing that, in all probability, the species is not closely circumscribed.

Cultures made by Dr. Fr. Bubák in 1901 from Bohemian material brought to light a biological form which only infected *Berula angustifolia*.<sup>26</sup>

In 1902 Dr. H. Klebahn attempted to repeat Dietel's cultures, and found that teliosporic material from the same immediate region, the exact locality having been changed and the rust destroyed, gave abundant aecia on *Pastinaca sativa*, but only slightly infected *Hippurus vulgaris*, and infected *Sium latifolium* and *Glaux maritima* not at all.<sup>27</sup> The year following he carried out more extensive cultures. Teliosporic material raised from aeciospores on *Pastinaca sativa*, infected both *Pastinaca* and *Berula angustifolia*. Teliosporic material from central Germany

<sup>24</sup> See Jour. Myc. 12:19. 1906.

<sup>25</sup> Hedwigia 29:149. 1890.

<sup>26</sup> Centr. Bakt. 9:226. 1902.

<sup>27</sup> Jahrb. Hamb. Wiss. Anst. 20:33. 1903.

infected both these hosts. Similar material from northern Germany infected only *Berula*. In 1904 aecia were raised in a similar way on *Oenanthe aquatica*.<sup>28</sup>

One of the earliest cultures with this pleophagic species was made by Mr. C. B. Plowright with English material, raising aecia on *Glaux maritima*, which belongs to the *Primulaceae*.<sup>29</sup>

A careful microscopic study of all these forms shows close agreement in morphological characters, and altogether there appears to be no reason to treat these cultural forms other than biological races of a single species. *Scirpus fluviatilis* is the American representative of the European *S. maritimus*, and by many authors is given the latter name. The writer has examined American collections that appear to belong to this species, as follows: on *S. fluviatilis* from Iowa, Illinois, Kansas, Nebraska, South Dakota, Ohio, and Wisconsin; on *Cicuta maculata* from Illinois, Iowa, and Colorado; on *Oenanthe californica* from central California, *Sium cicutaefolium* from Wisconsin, and on *Glaux maritima* from Montana and Wyoming.

18. GYMNOSPORANGIUM JUNIPERI-VIRGINIANAE Schw.—Galls from a tree of *Juniperus virginiana* near the laboratory, brought in by Mr. F. D. Kern, were used for sowing May 1 on *Malus coronaria*, *Sorbus americana*, and *Crataegus Pringlei*. Only the first gave infection, which showed abundant pycnia on May 14, but failed to develop aecia on account of injuries.<sup>25</sup>

19. GYMNOSPORANGIUM GLOBOSUM Farl.—Galls from a tree of *Juniperus virginiana*, found in the vicinity of Lafayette, Ind., were brought in by Mr. Guy W. Wilson, and used for making three sowings on *Crataegus Pringlei*, and one each on *Malus coronaria* and *Amelanchier* sp. All sowings on *Crataegus* gave abundance of pycnia, and one plant finally produced well formed aecia, the other plants not growing well. No infection was obtained on the other two hosts.

Similar material was also received from Dr. John A. Sheldon, of Morgantown, W. Va., and sown May 19 on *Crataegus Pringlei*, *Malus coronaria*, and *Sorbus americana*, all giving rise to an abundance of pycnia. As none of the plants grew well, only the sowing on *Sorbus* formed aecia, these finally reaching maturity and showing the characteristic structure of the species.

These results are parallel with, and confirm the work by Dr. Roland Thaxter, done some years ago.<sup>26</sup>

<sup>28</sup> Ztschr. Pfl.-Kr. 15:74. 1905.

<sup>29</sup> Gard. Chron. III. 7:682. 1890.

<sup>25</sup> For previous cultures see Jour. Myc. 12:13. 1906.

<sup>26</sup> Proc. Amer. Acad. Sci. 22:263. 1887; Bot. Gaz. 14:167. 1889.



The following three species have never before been tested by means of cultures, so far as the writer knows. Although few in number, they make an important addition to our knowledge of life histories:

I. MELAMPSORA LINI (Link) Desmaz.—For a number of years attempts have been made to obtain cultures of this cosmopolitan rust, and learn its full cycle of development. Many collections from different parts of the country, gathered at different times from November to April, have been tested, but with uniform failure to secure germination of the teliospores. The most numerous and promising collections were sent by Professor H. L. Bolley, of North Dakota, but equally in vain until one made the last day of April on cultivated flax, dug from under a snow bank, was received. This showed strong germination of the teliospores, and on May 4 was sown on *Linum Lewisii*, *Larix laricina*, and the day following on *Tsuga canadensis*, and *Arisaema triphyllum*. No clues were available, but judging from the willow, poplar, and some other species of the same genus, it was assumed that it might be heteroecious. Nevertheless, on May 16 pycnia began to appear on the flax, and on May 21 aecia. The next sowing was made May 18 on *Linum usitatissimum*, plants of which had not before been available, giving rise to pycnia May 26, and aecia May 30. Another equally successful sowing on the same host was made May 29, but exact record for the appearance of the sori was not kept.

The pycnia are small, pale, and inconspicuous, although numerous. They are globoid, subepidermal, and without ostiolar filaments. The aecia are of the caecoma form, that is, are without peridia. They are also rather pale, and not as prominent as the uredinia, for which, however, they might easily be mistaken. So far as I can learn they have never been collected, although probably common throughout the world. This may be due to their being inconspicuous, and quite as much to the earliness of their appearance.

The economic bearing of the discovery is to some extent obvious. Knowing the autoecious nature of the rust makes the destruction of old flax straw in flax growing regions a matter of moment, in order to lessen and retard the appearance of the rust in growing fields of flax. Upon reporting the first success to Professor Bolley he replied under date of May 23, 1906: "I am very much pleased to receive your letter, for with the information I am able to determine the source of a rust epidemic that we had in our flax breeding plots last year. I now find everywhere in the flax stubble plenty of volunteer flax with almost every plant more or less infected with spermogonia and aecidia." It will now be possible to intelligently devise methods for the practical control of flax rust.

2. *UROMYCES* ON *JUNCUS TENUIS*.—This rust, which has generally passed under the name of *Uromyces Junci*, is very common in the United States east of the Rocky Mountains from Canada to the Gulf of Mexico. It is often highly parasitized, so that the sori contain few or no well grown teliospores. A number of attempts to bring the rust under culture have failed because the teliospores could not be made to germinate. In 1902 a sowing was made on *Iris versicolor* without success.

Unparasitized material on *Juncus tenuis* Willd. was found near Lafayette, Ind., April 3, that proved to be viable, and was sown April 20 on *Lactuca canadensis*, *L. virosa*, *Senecio obovatus*, *Rudbeckia laciniata*, *Ambrosia trifida*, *A. artemisiaefolia*, and *Silphium perfoliatum*. Much to our delight pycnia were observed on the last host April 30, and were followed by a great abundance of aecia May 7. Another sowing was made May 4 on *S. perfoliatum* and *S. terebinthinaceum*, and the next day on *Parthenium integrifolium*. Only the first was infected, showing pycnia May 17, and May 31. The plant of *S. terebinthinaceum* was weak and did not grow well. Later sowings May 9 on *Polymnia canadensis*, and June 5 on *Parthenium integrifolium* and *Silphium terebinthinaceum*, gave no infection.

Another collection of teliospores on *Juncus tenuis* was sent by Dr. John L. Sheldon from Morgantown, W. Va., which was found close to *Houstonia caerulea* bearing aecia. This material was sown May 19 on *Houstonia caerulea*, *H. purpurea* and *Silphium perfoliatum*. No infection occurred on the *Houstonias*, but pycnia appeared on the *Silphium* May 29, and aecia June 4, both abundant and well formed.

Since obtaining the unequivocal cultural results, a morphological study has been made of the rust, which clearly demonstrates that it is specifically distinct from *Uromyces Junci* (Desm.) Tul. As it appears never to have been recognized as an autonomous species, the following name and description are submitted:

***Uromyces Silphii* (Syd.) nom. nov.** (*Accidium compositarium Silphii* Burr. Saccardo, Syll. Fung. 7:798. 1888; *Aecidium Silphii* Sydow, Uredineen 1546.)

O. Pycnia chiefly epiphyllous, in small groups, golden brown, subglobose or ellipsoid, subepidermal, 80-100  $\mu$  broad by 80-110  $\mu$  high; ostiolar filaments up to 65  $\mu$  long.

I. Aecia amphigenous, in groups 4-10 mm. across, crowded about the pycnia on discolored spots, deep-seated, short, 0.2-0.4 mm. in diameter; peridium colorless, margin recurved, lacerate, peridial cells rhombic, 20-28  $\mu$  across, outer wall thick, 6-8  $\mu$ , transversely striate, inner wall thinner, 4-6  $\mu$ , moderately verrucose; aeciospores angularly globoid, small, 13-18  $\mu$  in diameter, wall colorless, thin, about 1  $\mu$ , minutely verrucose.

On *Silphium integrifolium* Michx., Illinois (type, McLean County, May 3, 1881 or 1882, *A. B. Seymour* 4852, recorded in Burrill's Parasitic Fungi of Illinois, page 231), Wisconsin; *S. terebinthinaceum* Jacq., Illi-

nois, Wisconsin, Missouri; *S. perfoliatum* L., Indiana, Iowa, Wisconsin; *S. laciniatum* L., Illinois, Iowa, Kansas.

II. Uredinia amphigenous, scattered, roundish or somewhat elongated, small, 0.2-0.3 mm. wide, by 0.3-0.5 mm. long, tardily naked, dark cinnamon-brown, ruptured epidermis not conspicuous; urediniospores broadly ellipsoid or sometimes obovate, 13-19 by 15-23  $\mu$ , wall golden yellow, about 1.5  $\mu$  thick, sparsely and bluntly echinulate, pores 5 or 6, scattered.

III. Telia amphigenous, scattered, roundish or somewhat elongated, small, 0.2-0.3 mm. wide by 0.2-0.6 mm. long, tardily naked, firm, somewhat pulvinate, blackish brown, ruptured epidermis noticeable; teliospores angularly obovate, rounded, truncate or occasionally pointed above, usually narrowed below, 12-19 by 26-35  $\mu$ , wall chestnut-brown, 1.5-2  $\mu$  thick, much thicker above, 7-10  $\mu$ , smooth; pedicel light chestnut-brown, one to one and a half times length of spore.

On *Juncus tenuis* Willd., Indiana, Iowa, Wisconsin, Michigan, Minnesota, South Dakota, Missouri, New York, Maine, Massachusetts, West Virginia, Louisiana, Texas; *J. dichotomus* Ell., Florida.

It has been issued in the following exsiccati: aecial stage—Ellis & Ev. Fungi Columb. 1478; Sydow, Ured. 1546; telial stage—Seym. & Earle, Econ. Fungi 52, 528; Griffiths, W. Am. Fungi 244 (host *J. tenuis* not *J. longistylis*); Ellis & Ev. Fungi. Columb. 2394.

The following key will serve to separate the three common species of *Uromyces* on *Juncus*, when the urediniospores are present. In the absence of uredinia the urediniospores can usually be found to some extent in telial sori, even those that have withstood the winter, and are collected in the spring following their maturity, and especially so if they are parasitized.

Urediniospores verrucose, pores 2, equatorial.....*U. Junci* (Desm.) Tul.  
 Urediniospores echinulate, pores 4, equatorial.....*U. effusus* Arth.  
 Urediniospores echinulate, pores 5-6, scattered....*U. Silphii* (Syd.) Arth.

3. GYMNOSPORANGIUM NELSONI Arth. — At the time this species was published it was stated that Prof. Aven Nelson, the collector of the type material, considered it highly probable that the aecia found on *Amelanchier* in the vicinity belonged to the species. Teliosporic material on *Juniperus scopulorum* Sarg., sent by Mr. E. Bethel from Colorado this spring, gave the first opportunity to test the suggestion. Sowings were made May 29 on the leaves of *Amelanchier canadensis*, *Sorbus americana*, *Crataegus Pringlei*, *Pyrus japonica*, and *Aronia nigra*. On June 12 a few pycnia were observed on the *Amelanchier* and *Sorbus*, but owing doubtless to indifferent growth of the hosts no aecia were formed. The other hosts remained without infection. This helps in a small way to confirm Professor Nelson's suggestion, but is not conclusive.

#### SUMMARY.

The following is a complete list of successful cultures made during the season of 1906. It is divided into two series: species previously reported by the writer or other investigators, and species now reported for the first time.



*A. Species previously reported.*

1. MELAMPSORA BIGELOWII Thuem. — Teliospores on *Salix* sp. sown on *Larix decidua* Mill.

2. CRONARTIUM QUERCUS (Brond.) Schroet. — Aeciospores on *Pinus virginiana* Mill. sown on *Quercus velutina* Lam.

3. PUCCINIA OPIZII Bubák. — Teliospores on *Carex* sp. sown on *Lactuca canadensis* L., *L. virosa* L. and *L. sativa* L.

4. PUCCINIA SAMBUCI (Schw.) Arth. — Teliospores on *Carex Frankii* Kunth sown on *Sambucus canadensis* L.

5. PUCCINIA PECKII (DeT.) Kellerm. — Teliospores on *Carex trichocarpa* Muhl., and also on *C. lanuginosa* Michx., sown on *Onagra biennis* (L.) Scop.

6. PUCCINIA ALBIPERIDIA Arth. — Teliospores on *Carex squarrosa* L., *C. tetanica* Schk., and *C. crinita* Lam. sown on *Ribes rotundifolium* Michx., *R. gracile* Michx. and *R. cynosbati* L.

7. PUCCINIA ANGUSTATA Peck. — Teliospores on *Scirpus atrovirens* Muhl. sown on *Lycopus americanus* Muhl.

8. PUCCINIA ELEOCHARIDIS Arth. — Teliospores on *Eleocharis palustris* (L.) R. & S. sown on *Eupatorium perfoliatum* L.

9. PUCCINIA ANDROPOGONIS Schw. — Teliospores on *Andropogon scoparius* Michx. sown on *Pentstemon hirsutus* (L.) Willd.

10. PUCCINIA TOMIPARA Trel. — Teliospores on *Bromus purgans* L. sown on *Clematis virginiana* L.

11. PUCCINIA SUBNITENS Diet. — Teliospores on *Distichlis spicata* (L.) Greene sown on *Chenopodium album* L.

12. PUCCINIA POCULIFORMIS (Jacq.) Wettst. — Teliospores on *Agrostis alba* L. and on *Elymus canadensis* L. sown on *Berberis vulgaris* L., and aeciospores from *B. vulgaris* L. sown on *Hordeum vulgare* L. and *Triticum vulgare* Vill.

13. PUCCINIA TRANSFORMANS Ellis & Ev. — Teliospores on *Stenolobium Stans* (L.) Don sown on same host.

14. PUCCINIA XANTHII Schw. — Teliospores on *Xanthium* sp. sown on same host.

15. PUCCINIA SILPHII Schw. — Teliospores on *Silphium integrifolium* Michx. sown on *S. integrifolium* Michx., *S. perfoliatum* L., and *S. terebinthinaceum* Jacq.

16. PUCCINIA PRUNI-SPINOSAE Pers. — Aeciospores on *Hepatica acutiloba* DC. sown on *Prunus serotina* Ehrh. and *P. pumila* L.

17. UROMYCES SCIRPI (Cast.) Burr. — Teliospores on *Scirpus fluviatilis* (Torr.) A. Gray sown on *Cicuta maculata* L.

18. GYMNOSPORANGIUM JUNIPERI-VIRGINIANAE Schw. — Teliospores on *Juniperus virginiana* L. sown on *Malus coronaria* (L.) Mill.

19. GYMNOSPORANGIUM GLOBOSUM Farl. — Teliospores on *Juniperus virginiana* L. sown on *Crataegus Pringlei* Sarg., *Sorbus americana* Marsh., and *Malus coronaria* (L.) Mill.

*B. Species reported now for the first time.*

1. MELAMPSORA LINI (Link) Desmaz. — Teliospores on *Linum usitatissimum* L. sown on *L. Lewisii* Pursh and *L. usitatissimum* L.

2. UROMYCES SILPHII (Syd.) Arth. — Teliospores on *Juncus tenuis* Willd. sown on *Silphium perfoliatum* L.

3. GYMNOSPORANGIUM NELSONI Arth. — Teliospores on *Juniperus scopulorum* Sarg. sown on *Amelanchier canadensis* (L.) Medic. and *Sorbus americana* Marsh.

---

## AN HISTORICAL REVIEW OF THE PROPOSED GENERA OF PHYCOMYCETES.

### I. PERONOSPORALES.

GUY WEST WILSON.

In the present consideration of the generic types of the *Phycomycetes* the genera will be arranged chronologically under each order, with the type species, the synonyms, the homonyms, and such other information under each genus as may seem desirable. This is followed by an alphabetical list of the genera with their type species, in which the tenable names are printed in black type while those which are untenable are in common type.

The subject of the generic types of the *Phycomycetes* was first taken up at the suggestion of Dr. J. C. Arthur, while a student in his laboratory, and the results embodied in a thesis which was presented to the Faculty of Purdue University, to the authorities of which institution I am indebted for permission to publish the material contained in the thesis. I wish to also express my hearty appreciation of the courtesies shown me by Dr. Arthur and by Dr. J. H. Barnhart in the way of critical and bibliographical assistance and the loan of otherwise inaccessible books and by those in charge of the various libraries consulted both in Lafayette and New York.



## 1. ALBUGO Roussel, Fl. Calvados, ed. 2. 47. 1806.

The name *Albugo* was first used for a subgenus by Persoon\* who included under it the white spored species of *Uredo*, of which he recognized two, *U. candida* and *U. Cheiranthi*, the first of which is the type species. Roussel elevated the subgenus to generic rank with Persoon's first species as the monotype of the genus. In this species he followed Persoon in including the white rust of the *Cruciferae* and of *Tragopogon*, and by way of good measure added some fungus on *Urtica* which in all probability belongs to the *Uredinales*. The genus is usually credited to S. F. Gray† who used the name in the same sense fifteen years later.

Synonym: *Cystopus* Lév., not Blume.

2. PERONOSPORA<sup>1</sup> Corda, Icon. Fung. 1: 20. 1837.

This genus was published for a single species, *P. rumicis*, which is figured on pl. 5. f. 273.

## 3. BREMIA Regel, Bot. Zeit. 1:666. 1843.

The only species which has so far been referred to the genus is *B. Lactucae* Regel, a fairly good figure of which (pl. 3B) accompanies the description.

Synonym: *Actinobotrys* Hoffm.

## 4. CYSTOPUS Léville, Ann. Sc. Nat. III. 8: 371. 1847.

The genus is characterized and five species of *Uredo*, of which the first is *U. candida*, are cited as members of the genus. Except the last species named these are all congeneric. While it is customary to credit to Léville the names of all these species under the genus *Cystopus*, he refrained, not only in this but in subsequent papers, from forming the combinations with which he is so generously credited by his contemporaries. The name is untenable as the genus is a typonym of *Albugo*. It has also been applied to an *Orchidaceous*‡ genus.

## 5. ACTINOBOTRYS Hoffman, Bot. Zeit. 14: 154. 1856.

But one species, *A. Tulasnei*, was described, the figure of a portion of the conidiophore of which (pl. 5. f. 22) is unmistakably that of *Bremia Lactucae* Regel, 1843.

6. BASIDIOPHORA<sup>1</sup> Roze & Cornu, Ann. Sci. Nat. V. 11: 84. 1869.

This genus is founded on a single species, *B. entospora*, which is described at great length and carefully figured (pl. 4).

Synonym: *Gilletia* Sacc. & Penz.

\* Syn. Meth. Fung. 223. 1801.

† Nat. Arr. Brit. Pl. 1:540. 1821.

‡ *Cystopus* Blume, Orch. Archip. Ind. 82. 1858.

7. PHYTOPHTHORA de Bary, Jour. Roy. Agr. Soc. II. 12: 240. 1876.

The present genus was founded for a single species, *Peronospora infestans*, which is renamed *Phytophthora infestans* de Bary. The species and its life history are discussed in detail and illustrations are given (f. 3, 4). A duly accredited reprint of this article§ is usually cited instead of the original place of publication.

8. SCLEROSPORA Schröter, de Bary, Bot. Zeit 39: 621. 1881.

The present name was used by Schröter\* as a subgenus of *Peronospora*, under which there was placed a single species previously described by Saccardo as *Protomyces graminicola*. The first use of the name in a generic sense is in a paper by de Bary in which he enumerates the valid genera of *Peronosporaceae* among which is *Sclerospora* Schröter. The first binominal combined with the generic name is in Cohn's *Kryptogamen Flora von Schlesien*|| where the type of the subgenus of 1879 becomes *S. graminicola* (Sacc.) Schröt. Up to this time no other species had been associated with the name *Sclerospora* whether regarded as a genus or as a subgenus.

9. GILLETIA Saccardo & Penzig; Sacc. Michelia 2: 587. 1882.

The only species which is referred to this genus is *G. spinuligera* Sacc. & Therry, of which the authors say "Oogonia ignota, sed totus fungi habitus peronosperoides." The fungus is co-specific with *Basidiophora entospora* Roze & Cornu. The type specimens of both these species were collected in France on *Leptilon canadensis* (L.) Britton. The same generic name has since been used in *Commelinaceae*.¶

10. PLASMOPARA Schröter, in Cohn, Krypfl. Fl. Schles. 3: 236. 1886.

Eight species were referred to this genus by its author who founded it as "Gatt. **Plasmopara**. (*Peronospora* Sect. I. *Zoosporiparac* und Sect. II. *Plasmatoparac* de Bary)." The generic name is merely a modification of de Bary's second sectional name, which implies that the conidia germinate by a plasma. This is true only of de Bary's second section, the species of the first germinating by zoospores. Two species, *Peronospora pygmaea* Unger and *P. densa* Rabenh., are included by de Bary in his section *Plasmatoparac* of which *P. pygmaea* is the type. As

§ Jour. Bot. 14:105-126, 148-154. 1876.

\* Hedwigia, 18:86. 1879.

|| 3<sup>1</sup>:236. 1886.

¶ Rendle, Jour. Bot. 84:55. 1896.

this species is the first one from this section which is cited by Schröter it is the type of the genus. Of his other species two were known to de Bary, while the others, including the type of *Basidiophora*, are of later date.

11. CHLOROSPORA Spegazzini, Rev. Argent. Hist. Nat. 1: 29. 1891.

The single species of this genus, *C. vastatrix*, is described as having monopodially branched conidiophores and colored conidia which germinate by a plasma. The genus is, therefore, very close to *Plasmopora*.

12. DREPANOCONIS Schröt. & P. Henn.; Henning's, Hedwigia 35: 211. 1896.

This genus, which is placed by its authors in *Albuginaceae*, contains a single species *D. brasiliensis*, the true position of which is probably among the *Hyphomycetes*.

13. PSEUDOPERONOSPORA Rostewzew, Ann. Inst. Agron. Moscow 9:47. 1903; Flora 92:422. 1903.

The present genus was founded upon *Peronospora cubensis* B. & C. and a Russian variety of that species. The characters of these fungi and their relatives are discussed in detail and the article, which is in Russian, is profusely illustrated. There is no room to question the identity of the genus and its type. A translation of the article appeared in Flora about eight or ten months later.

Synonym: *Peronoplasmopora* (Berlese) Clinton.

14. KAWAKAMIA Miyabe, Mag. Bot. Tokyo 17:(306). 1903.

This genus is founded on a single species, *Peronospora cypripetidis* Miyabe & Ideta. Although published in the Japanese section of the Magazine the descriptions of both genus and species are in English.

15. PERONOPLASMOPARA Clinton, Ann. Rep. Conn. State Agr. Exp. St. 29:234. 1905.

Berlese\* proposed this name for a subgenus of *Plasmopora* in which he included two species, *Plasmopora Celtidis* (Waite) Berlese and *P. cubensis* (B. & C.) Humphrey. He described both species and figured the latter, which is therefore the subgeneric type. When Clinton elevated the subgenus to generic rank he designated "Types: *Peronoplasmopora cubensis* (B. & C.) Clinton, *Peronoplasmopora celtidis* (Waite) Clinton." As it is manifestly impossible for a genus to have more than one type, and as his discussion is based almost entirely upon the first species which he mentions we must consider both subgenus and genus as founded upon *Peronospora cubensis* B. & C.

\* Riv. Pat. Veg. 9:123. 1900.



16. PHLEOPHYTHORA Klebahn, Cent. Bakt II. 15:336. 1905.

The genus is founded upon an imperfectly known species, *P. syringae*, which may belong either to the *Peronosporales* or to the *Anclystidales*.

17. MYCELOPHAGUS Mangin, Compt. Rend. Acad. Sci. Paris 136:472. 1906.

The genus is founded on a poorly described and imperfectly known species, *M. castaneae*, which probably belongs to *Anclystidaceae*.

### Alphabetical List of Genera.

(The tenable names in black type.)

*Actinobotrys* Hoffm. 1856.—*A. Tulasnei* Hoffm. = *Bremia lactucae* Regel. 1843.

*Albugo* (Pers.) Rouss. 1806.—*A. candida* (Pers.) Rouss.

*Basidiophora* Roze & Cornu, 1869.—*B. entospora* Roze & Cornu.

*Bremia* Regel, 1843.—*B. lactucae* Regel.

*Chlorospora* Speg. 1891.—*C. vastatrix* Speg.

*Cystopus* Lév. 1847.—*C. candidus* (Pers.) Lév. 1848. = *Albugo candida* Rouss. 1806.

*Drepanoconis* Schröt. & P. Henn. 1896.—*D. brasiliensis* Schröt. & P. Henn. Hyphomycete?

*Gilletia* Sacc. & Penz. 1882.—*G. spinuligera* Sacc. & Therry. = *Basidiophora entospora* Roze & Cornu. 1869.

*Kawakamia* Miyabe, 1903.—*K. cyperi* (Miyabe & Ideta) Miyabe.

*Mycelophagus* Mangin, 1906.—*M. castaneae* Mangin. Tenable as a genus but probably belonging to another order of Oömycetes.

*Peronoplasmopara* (Berlese) Clinton, 1905.—*P. cubensis* (B. & C.) Clinton. = *Pseudoperonospora cubensis* (B. & C.) Rostew. 1903.

*Peronospora* Corda, 1837.—*P. rumicis*. Corda.

*Phleophythora* Klebahn, 1905.—*P. syringae* Klebahn.

*Phytophthora* de Bary, 1876.—*P. infestans* (Mont.) de Bary.

*Plasmopara* Schröt. 1886.—*P. pygmaca* (Unger) Schröt.

*Pseudoperonospora* Rostew. 1903.—*P. cubensis* (B. & C.) Rostew.

*Sclerospora* (Schröt.) de Bary, 1881.—*S. graminicola* (Sacc.) Schröt. 1886.

## INDEX TO NORTH AMERICAN MYCOLOGY.

*Alphabetical List of Articles, Authors, Subjects, New Species and Hosts, New Names and Synonyms.*

W. A. KELLERMAN.

ADDITIONS, Some, To the Flora of Middlesex Co., Massachusetts. [24 species of lichens.] Reginald Heber Howe Jr. Bryologist 9:81-2. Sept. 1906.

AECIDIUM cannonii Griffiths n. sp. on Fouquieria splendens Engelm. Bull. Torr. Bot. Club, 34:210. April 1907.

AECIDIUM ludwigiae E. & E., syn. of Puccinia ludwigiae q. v.

AECIDIUM sarcobati Peck on Nitrophila occidentalis. [Griffiths.] Bull. Torr. Bot. Club, 34:210. April 1907.

AEGERITA candida Pers., the conidial form of Peniophora candida (Pers.) Lyman n. sp. Proc. Boston Soc. Nat. Hist. 33:168. February 1907.

AGARICUS retirugis Fries, [syn. of Panaeolus carbonarius Batsch. Morgan]. Jour. Mycol. 13:61. Mar. 1907.

ALBOCOCCUS Winslow & Rogers n. gen. Coccaceae. Science N. S. 25:815. 24 May 1907.

ALLODUS nesaeae Arth., syn. of Puccinia ludwigiae q. v.

ARTHUR, J. C. McAlpine's Studies of Australian Rusts [review]. Jour. Mycol. 13:41-2. Mar. 1907.

ASCOGENOUS forms of Gloeosporium and Colletotrichum. C. L. Shear and Anna K. Wood. Bot. Gaz. 43:259-266. April 1907.

ASCOMYCETES and Lower Fungi [notes on exsiccati, 25 nos.] Guy West Wilson and Fred Jay Seaver. Jour. Mycol. 13:48-52. Mar. 1907.

AURISCALPIUM auriscalpium (L.) S. F. Gray. [Hydnum auriscalpium L.; Auriscalpium vulgare Karsten; Pleurodon auriscalpium Karsten; Leptodon auriscalpium Quélet. Mem. Torr. Bot. Club, 12:178. 13 June 1906.

AURISCALPIUM vulgare Karsten, syn. of Auriscalpium auriscalpium q. v.

AUROCOCCLUS Winslow & Rogers n. gen. Coccaceae. Science N. S. 25:815. 24 May 1907.

BACTERIOLOGISTS, Society of, 8th Annual Meeting [titles and abstracts of papers]. C. S. Prescott, Secretary. Science N. S. 25:805-820. May 24 1907.

- BACTERIUM tumefaciens Smith & Townsend n. sp., a Schizomycete causing rapid multiplication of the young tissues of *Chrysanthemum frutescens*, *Prunus persica*, etc., [tumors]. Science N. S. 25:671-3. 26 April, 1907.
- BANKER, Howard James. A contribution to a Revision of the North American Hydnaceae. Mem. Torr. Bot. Club, 12: 99-194. 13 June 1906.
- BARTHOLOMEW, Elam Fungi Columbiani, Century XXIII. Dec. 10, 1906.
- BARTHOLOMEW, Elam. Fungi Columbiani, Century XXIV. March 15, 1907.
- BERGEN, Joseph Y., and Davis, Bradley M. Type-studies — Schizomycetes, Saccharomycetes, Phycomycetes, Ascomycetes. Lab. & Field Man. Bot. 102-117. 1907.
- BERMUDA grass. See *Cynodon dactylon*.
- BESSEY, Ernst A. Spore forms of *Spegazzinia ornata* Sacc. Jour. Mycol. 13:43-5. Pl. 101. Mar. 1907.
- BIRD'S-NEST fungi. C. G. Lloyd's Myc. Notes. No. 24, Dec. 1906, pp. 293-308 pl. 91-93.
- BLAKESLEE A. F. Heterothallism in Bread Mould. *Rhizopus nigricans*. Bot. Gaz. 43:415-418. June 1907.
- BOISDUVALIA glabella (Nutt.) Walp. host to *Puccinia glabella* Holway n. sp. Holway's N. A. Uredin. 1:76. 10 May 1907.
- BURLINGHAM, Gertrude Simmons. Suggestions for the study of Lactariae. Torreya, 7:118-123. June 1907.
- CALIFORNIA Lichens. Lichen Distribution in the Santa Cruz Peninsula, California. Bot. Gaz. 43:267-273. April 1907.
- CALODON aurantiacus Karst., syn. of *Hydnellum floriforme* q. v.
- CALODON cyathiformis Quélet, syn. of *Phellodon tomentosus* q. v.
- CALODON ferrugineus Karst., syn. of *Hydnellum sanguinarium* q. v.
- CALODON niger Quélet, syn. of *Phellodon niger* q. v.
- CALODON suaveolens, Karst., syn. of *Hydnellum suaveolens* q. v.
- CALODON velutinus Karst., syn. of *Hydnellum velutinum* q. v.
- CALODON zonatus Karst., syn. of *Hydnellum zonatum* q. v.
- CAMPBELL, Douglas Houghton. Fungi. A University Text-book of Botany. 2d ed. 149-193. 1907.
- CERATOSTOMA biparasiticum E. & E. syn. of *Melanospora parasitica* q. v.



- CERATOSTOMELLA capillifera Hedgcock n. sp. on lumber made from Liquidamber styraciflua L. An. Rep. Mo. Bot. Gar. 1906, 17:71. 27 Sept. 1906.
- CERATOSTOMELLA minor Hedgcock n. sp. An. Rep. Mo. Bot. Gar. 1906, 17:74. 27 Sept. 1906.
- CERATOSTOMELLA pluriannulata Hedgcock n. sp. on a block of blued sapwood from Quercus rubra. An. Rep. Mo. Bot. Gar. 1906, 17:72. 27 Sept. 1906.
- CERATOSTOMELLA schrenkiana Hedgcock n. sp. on sapwood of pine lumber made from Pinus echinata. An. Rep. Mo. Bot. Gar. 1906, 17:67. 27 Sept. 1906.
- CLADONIAS, Further notes on, VIII. Bruce Fink. Bryologist, 9:89-91 Pl. VIII. Nov. 1906.
- CLADONIAS, Further Notes on, XI. Cladonia pyxidata and Cladonia pityrea. Bruce Fink. Bryologist, 10:57-60 Pl. VIII. July 1907.
- CLEMATIS paniculata, cult., host to Diplodia hortensis Sacc. n. sp. Jour. Mycol. 13:47. Mar. 1907.
- CLIMACODON septentrionalis Karsten, syn. of *Steccherinum septentrionale* q. v.
- CLIMACODON ochraceus Karst., syn. of *Steccherinum ochraceum* q. v.
- CLINTON, George Perkins. Ustilaginales (Ustilaginaceae and Tilletiaceae) [Monograph]. North American Flora 7:1-82. 4 Oct. 1906.
- CHLORIS elegans H. B. K., host to Sclerospora farlowii Griffiths n. sp. Bull. Torr. Bot. Club, 34:207. April 1907.
- CHROMOGENIC FUNGI, Studies upon some which discolor wood. George Grant Hedgcock. An. Rep. Mo. Bot. Gar. 1906, 17:59-114. Pl. 3-11. 27 Sept. 1906.
- COCCACEAE, Generic characters in the. C. E. A. Winslow and Anne F. Rogers. [Abstract.] Science N. S. 25:813-4. 24 May 1907.
- COLLETOTRICHUM, Gloeosporium and, Ascogenous forms of. C. L. Shear and Anna K. Wood. Bot. Gaz. 43:259-266. April 1907.
- CONCERNING Some West American Fungi [several new species]. David Griffiths. Bull. Torr. Bot. Club, 34:207-211. April 1907.
- CORTICIUM subiganteum Berk. [Organic connection between this species and Michenera artocreas B. & C., the imperfect form, Lyman]. Proc. Boston Soc. Nat. Hist. 33:151. Feb. 1907.

- CORTINARIUS rubipes Kauffman n. sp.; the stem is attached to roots of *Acer saccharinum* and *Quercus rubrum* on which it forms mycorrhiza. Mich. Acad. Sci. 8:32. Meeting of 1906. Date (?).
- COSMOS sulphureus, host to *Entyloma holwayi* Sydow. [Mexico.] N. A. Flora, 7:63. 4 Oct. 1906.
- CRATAEGUS, branch, host to *Dermatea olivascens* Rehm n. sp. Ann. mycolog. 5:80. Feb. [issued 30 Mar.] 1907.
- CULTURE Studies on Polymorphism of Hymenomycetes. George Richard Lyman. (Proc. Boston Soc. Nat. Hist. 33:125-209. Pl. 18-26. Feb. 1907.
- CYNODON dactylon Pers., Bermuda grass, host to *Ustilago cynodontis* P. Henn. Bull. Torr. Bot. Club, 34:208. April 1907.
- DANDENO, J. B. A fungus disease of greenhouse Lettuce, [*Didymaria perforans*]. Mich. Acad. Sci. 1906, 8:45-7. Date of issue?
- DAVIS, Bradley M., see *Bergen, Joseph Y. and . . .*
- DERMATEA olivascens Rehm. n. sp. on (dead?) branches of *Crataegus* still on the tree. Ann. Mycolog. 5:80. Feb. 1907. [Issued 30 Mar. 1907.]
- DICAEOMA nesaeae Kuntze, syn. of *Puccinia ludwigiae* q. v.
- DIPLODIA hortensis Sacc. n. sp. in caulibus *Clematidis paniculatae*. Jour. Mycol. 13:47. Mar. 1907.
- DIRINA franciscana A. Zahlbruckner n. sp., on rocks 50 to 75 feet above the sea, San Francisco. Bot. Gaz. 43:270. April 1907.
- DISEASES, Plant, in 1906. W. A. Orton. Yearbook U. S. Dept. Agr. 1906:409-508. 1907.
- DRYODON coralloides Karst., syn. of *Hericium coralloides* q. v.
- DRYODON erinaceus Quél., syn. of *Hericium erinaceus*.
- ECHINODONTIUM tinctorium E. & E. [*Fomes tinctorius* E. & E.; *Hydnum tinctorium* Lloyd; *Hydnofomes tsugicola* Hennings & Shirai]. Mem. Torr. Bot. Club, 12:136. 13 June 1906.
- ENTYLOMA holwayi Sydow, on *Cosmos sulphureus*. [Mexico.] N. A. Flora, 7:63. 4 Oct. 1906.
- ERYSIPHOSIS Halst. [a gen. *Erysiphe* non differt immo *E. parnassiae* est forma *Erysiphe polygoni*. Salmon.] Sacc. Syll. 16:399. 1 Feb. 1902.
- ?EUCHLAENIA luxurians, host to *Ustilago kellermanii* Clinton n. sp. [Guatemala.] N. A. Flora, 7:15. 4 Oct. 1906.

- EUROTIA lanata Moq., host to Puccinia eurotiae Griffiths n. sp. Bull. Torr. Bot. Club, 34:211. April 1907.
- FAWCETT, Edna H., see *Kellerman, Karl F.* and . . .
- FAWCETT, H. S. Report of Assistant in Botany and Horticulture. [Diseases of Plants.] Fla. Agr. Exp. Sta. An. Report 1906:XXI-XXVII. 1907.
- FOUQUIERIA splendens Engelm., host to Aecidium cannonii Griffiths n. sp. Bull. Torr. Bot. Club, 34:210. April 1907.
- FINK, Bruce. Further notes on Cladonias. VIII. Bryologist, 9:89-91. Pl. VIII. Nov. 1906.
- FINK, Bruce. Cladonias, Further Notes on, XI. Cladonia pyxidata and Cladonia pityrea. Bryologist, 10:57-60 Pl. VIII. July 1907.
- FOLIACEOUS and fruticous Lichens, A list of, collected at Chilson Lake, Essex Co., New York. Altitude 1200 feet. Carolyn W. Harris. Bryologist, 9:48-52. May 1906.
- FOMES tinctorius E. & E., syn. of *Echinodontium tinctorium* q. v.
- FORM-STIMULUS, Gravity as a, in fungi. Heinrich Hasselbring. Bot. Gaz. 43:251-8. April 1907.
- FRIESITES caput-ursi Karsten, syn. of *Hericium caput-ursi* q. v.
- FRIESITES coralloides Karst., syn. of *Hericium coralloides* q. v.
- FRUTICOUS Lichens, see *Foliaceous and fruticous, A list of. Essex Co., New York.* . . .
- FUNGI. In Douglas Houghton Campbell's University Text-book of Botany. 2d ed. 149-193. 1907.
- FUNGI Columbiani, Century XXIII. Elam Bartholomew, Dec. 10, 1906.
- FUNGI Columbiani, Century XXIV. Elam Bartholomew, March 15, 1907.
- FUNGI, Concerning Some West American [several new species]. David Griffiths. Bull. Torr. Bot. Club, 34:207-211. April 1907.
- FUNGI Utahenses, Fascicle five. [Exsiccati Nos. 101-125.] A. O. Garrett. 19 Jan. 1907.
- FUNGUS Disease, A, of greenhouse Lettuce [*Didymaria perforans*]. J. B. Dandeno. Mich. Acad. Sci. 1906, 8:45-7. Date of issue?
- GARRETT, A. O. Fungi Utahenses, Fascicle five. [Exsiccati Nos. 101-125.] 19 Jan. 1907.



- GENERIC characters in the Coccaceae. C. E. A. Winslow and Anne F. Rogers. [Abstract.] *Science*, N. S. 25:813-4. 24 May 1907.
- GLOEOSPORIUM and Colletotrichum. C. L. Shear and Anna K. Wood. *Bot. Gaz.* 43:259-266. April 1907.
- GLOIODON strigosus Karsten, *syn. of Steccherinum strigosum* q. v.
- GLONIOPSIS smilacis (Schw.) Wilson & Seaver n. n. [Hysterium smilacis Schw.; Hypoderma smilacis (Schw.) Sacc.; Hysterographium smilacis (Schw.) E. & E.] *Jour. Mycol.* 13:49. Mar. 1907.
- GRANDINIODES Banker n. g. Hydnaceae. *Mem. Torr. Bot. Club*, 12:179. 13 June 1906.
- GRANDINIODES flavum (Swartz) Banker n. n. [Peziza flava Swartz; Hydnum flavum Berkeley; Hydnum brunneo-leucum B. & C.] *Mem. Torr. Bot. Club*, 12:179. 13 June 1906.
- GRASS-KILLING Slime Mould. [Physarum cinereum Pers.] John W. Harshberger. *Proc. Am. Philosophical Soc. Philadelphia*, 45:271-3. Oct.-Dec. 1906.
- GRAVITY as a Form-stimulus in Fungi. Heinrich Hasselbring. *Bot. Gaz.* 43:251-8. April 1907.
- GRIFFITHS, David. Concerning some West American Fungi [several new species]. *Bull. Torr. Bot. Club*, 34:207-211. April 1907.
- HARRIS, Carolyn W. Lichens of the Adirondack League Club Tract [list of 60 species]. *Bryologist*, 10:64-6. July 1907.
- HARRIS, Carolyn W. A list of foliaceous and fruticous lichens collected at Chilson Lake, Essex Co., New York. Altitude 1200 feet. *Bryologist*, 9:48-52. May 1906.
- HARSHBERGER, John W. A Grass-killing Slime Mould. *Physarum cinereum*. *Proc. Am. Philosophical Soc. Philadelphia*, 45:271-3. Oct.-Dec. 1906.
- HASSELBRING, Heinrich. Gravity as a Form-stimulus in fungi. *Bot. Gaz.* 43:251-8. April 1907.
- HEDGCOCK, George Grant. Studies upon some Chromogenic Fungi which discolor wood. *An. Rep. Mo. Bot. Gar.* 1906, 17:59-114. Pl. 3-11. 27 Sept. 1906.
- HEDGCOCK, George Grant. Zonation in Artificial Cultures of Cephalothecium and other Fungi. *An. Rep. Mo. Bot. Gar.* 1906, 17:115-7. Pl. 13-16. 27 Sept. 1906.
- HERICIUM, Key to species. [Banker.] *Mem. Torr. Bot. Club*, 12:113. 13 June, 1906.

- HERICIUM* caput-ursi (Fries) Banker n. n. [*Hydnum* caput-ursi Fries; *Friesites* caput-ursi Karsten; *H. caput-ursi brevispirem* Peck.] Mem. Torr. Bot. Club, 12:118. 13 June 1906.
- HERICIUM* coralloides (Scop.) Pers. [*Hydnum* coralloides Scop.; *H. crispum* Scop.; *Medusina coralloides* Chev.; *Dryodon coralloides* Karst.; *Friesites coralloides* Karst.] Mem. Torr. Bot. Club, 12:115. 13 June 1906.
- HERICIUM* croceum (Schw.) Banker n. n. [*Sistotrema croceum* Schw.; *Hydnum croceum* Schw.; *H. scheidermayeri* Heuf.] Mem. Torr. Bot. Club, 12:121. 13 June 1906.
- HERICIUM* erinaceus (Bull.) Pers. [*Hydnum* erinaceus Bull.; *Manina cordiformis* Scop.; *Medusina patula* Chev.; *Dryodon erinaceus* Quélet.] Mem. Torr. Bot. Club, 12:119. 13 June 1906.
- HERICIUM* fasciculare (Alb. & Schw.) Banker n. n. [*Hydnum fasciculare* A. & S.] Mem. Torr. Bot. Club, 12:123. 13 June 1906.
- HERICIUM* fimbriatum Banker n. sp. on a decaying stump of some hardwood, between the bark and the wood. Mem. Torr. Bot. Club, 12:122. 13 June 1907.
- HERICIUM* laciniatum [Leers] Banker n. n. [*Hydnum laciniatum* Leers; *H. ramosum* Bull.; *H. abietinum* Schrad.; *Medusina coralloides* Chev.] Mem. Torr. Bot. Club, 12:114. 13 June 1906.
- HERRE, Albert W. C. T. Lichen Distribution in the Santa Cruz Peninsula, California. Bot. Gaz. 43:267-273. April 1907.
- HETEROTHALLISM in Bread Mould, *Rhizopus nigricans*. A. F. Blakeslee. Bot. Gaz. 43:415-418. June 1907.
- HOWE, Jr., Reginald Heber. Some Additions to the Flora of Middlesex Co., Massachusetts. [24 species of lichens.] Bryologist 9:81-2. Sept. 1906.
- HOLWAY, E. W. D. North American Uredineae, Vol. I, Part III, pp. 57-80, pl. 24-34. 10 May 1907.
- HOLWAY, E. W. D. North American Uredineae. Part II, 1:53-56. Pl. 11-23. 15 May 1906.
- HOWE, Jr., Reginald Heber. Some Lichens of Mt. Watatic, Massachusetts. [List of Species.] Bryologist, 9:46-48. May 1906.
- HYDNACEAE, A Contribution to a Revision of the North American. Howard James Banker. Mem. Torr. Bot. Club, 12:99-194. 13 June 1906.

- HYDNACEAE, Synopsis of Genera. [Banker.] Mem. Torr. Bot. Club, 12:103. 13 June 1906.
- HYDNELLUM, Key to species. [Banker.] Mem. Torr. Bot. Club, 12:150-1. 13 June 1906.
- HYDNELLUM aurantiacum Karst., *syn. of Hydnellum floriforme q. v.*
- HYDNELLUM carbunculus (Secr.) Banker n. n. [Hydnum carbunculus Secr.] Mem. Torr. Bot. Club, 12:151. 13 June 1907.
- HYDNELLUM complicatum Banker n. sp. Mem. Torr. Bot. Club, 12:161. 13 June 1906.
- HYDNELLUM conigenum (Peck) Banker n. n. [Hydnum conigenum Peck.] Mem. Torr. Bot. Club, 12:160. 13 June 1906.
- HYDNELLUM concrescens (Pers.) Banker n. n. [Hydnum concrescens Pers.] Mem. Torr. Bot. Club, 12:157. 13 June 1906.
- HYDNELLUM cyaneotinctum (Peck) Banker n. n. [Hydnum cyaneotinctum Peck.] Mem. Torr. Bot. Club, 12:164. 13 June 1906.
- HYDNELLUM cyathiforme Karsten, *syn. of Phellodon tomentosus q. v.*
- HYDNELLUM earlianum Banker n. sp. on earth in woods. Mem. Torr. Bot. Club, 12:161. 13 June 1906.
- HYDNELLUM ferrugineum Karst. *syn. of Hydnellum sanguinarium q. v.*
- HYDNELLUM floriforme (Schaeff.) Banker n. n. [Hydnum floriforme Schaeff.; Hydnum suberosum aurantiacum Batsch; Hydnum compactum Pers.; Hydnum hybridum Pers.; Hydnum aurantiacum A. & S.; Hydnum aurantium Raf.; Hydnellum auranticum Karst.; Calodon aurantiacus Karst.; Phaeodon aurantiacus Schroet.] Mem. Torr. Bot. Club, 12:159. 13 June 1906.
- HYDNELLUM humidum (Banker) n. n. [Hydnum humidum Banker.] Mem. Torr. Bot. Club, 12:162. 13 June 1906.
- HYDNELLUM nigrum Karsten, *syn. of Phellodon niger q. v.*
- HYDNELLUM nuttallii Banker n. sp. on ground in woods. Mem. Torr. Bot. Club, 12:155. 13 June 1906.
- HYDNELLUM sanguinarium Banker n. n. [Hydnum ferrugineum Fries non Pers.; Phaeodon ferrugineus Schroet.; Calodon ferrugineus Karst.; Hydnellum ferrugineum Karst.] Mem. Torr. Bot. Club, 12:152. 13 June 1906.



- HYDNELLUM scrobiculatum (Fr.) Banker n. n. [Hydnum scrobiculatum Fr.] Mem. Torr. Bot. Club, 12:156. 13 June 1906.
- HYDNELLUM suaveolens (Scop.) Karst. [Hydnum suaveolens Scop.; Hydnum boreale Banker; Calodon suaveolens Karst.] Mem. Torr. Bot. Club, 12:162. 13 June 1906.
- HYDNELLUM velutinum (Fr.) Banker n. n. [Hydnum velutinum Fr.; Hydnum spongiosipes Peck; Calodon velutinus Karst.] Mem. Torr. Bot. Club, 12:153. 13 June 1906.
- HYDNUM zonatum (Batsch) Karst. [Hydnum zonatum Batsch; Calodon zonatus Karst.; Phaeodon zonatus Schroet.] Mem. Torr. Bot. Club, 12:158. 13 June 1906.
- HYDNOFOMES tsugicola Hennings & Shirai, *syn. of Echinodontium tinctorium q. v.*
- HYDNUM abietinum Schrad., *syn. of Hericium laciniatum q. v.*
- HYDNUM abietinum Schrad., *syn. of Hericium laciniatum q. v.*
- HYDNUM adustum Schw., *syn. of Steccherinum adustum q. v.*
- HYDNUM agaricoides Swartz, *syn. of Steccherinum agaricoides q. v.*
- HYDNUM albonigrum Peck, *syn. of Phellodon alboniger q. v.*
- HYDNUM atroviride Morgan, *syn. of Sarcodon atroviride q. v.*
- HYDNUM aurantiacum A. & S., *syn. of Hydnellum floriforme q. v.*
- HYDNUM aurantium Raf., *syn. of Hydnellum floriforme q. v.*
- HYDNUM auriscalpium L., *syn. of Auriscalpium auriscalpium q. v.*
- HYDNUM blackfordae Peck, *syn. of Sarcodon blackfordae q. v.*
- HYDNUM boreale Banker, *syn. of Hydnellum suaveolens q. v.*
- HYDNUM brunneo-leucum B. & C., *syn. of Grandinioides flazum q. v.*
- HYDNUM caput-ursi Fries, *syn. of Hericium caput-ursi q. v.*
- HYDNUM caput-ursi brevispinum Peck, *syn. of Hericium caput-ursi q. v.*
- HYDNUM carbunculus Secr., *syn. of Hydnellum carbunculus q. v.*
- HYDNUM cervinum Pers., *syn. of Sarcodon imbricatus q. v.*
- HYDNUM compactum Pers., *syn. of Hydnellum floriforme q. v.*
- HYDNUM conchiforme Sacc., *syn. of Steccherinum ochraceum q. v.*
- HYDNUM conigenum Peck, *syn. of Hydnellum conigenum q. v.*

- HYDNUM coraceo-membranaceum Schw., *syn. of Phellodon coraceo-membranaceus q. v.*
- HYDNUM concrescens Pers., *syn. of Hydnellum concrescens q. v.*
- HYDNUM coralloides Scop., *syn. of Hericium coralloides q. v.*
- HYDNUM croceum Schw., *syn. of Hericium coralloides q. v.*
- HYDNUM crispum Scop., *syn. of Hericium coralloides q. v.*
- HYDNUM cristatum Bres., *syn. of Sarcodon cristatus q. v.*
- HYDNUM cyaneotinctum Peck, *syn. of Hydnellum cyaneotinctum q. v.*
- HYDNUM cyathiforme Schaeffer non Bull., *syn. of Phellodon tomentosus q. v.*
- HYDNUM daviesii Sowerb., *syn. of Steccherinum ochraceum q. v.*
- HYDNUM delicatum Schw. non Klotsch, *syn. of Phellodon delicatus q. v.*
- HYDNUM erinaceus Bull., *syn. of Hericium erinaceus q. v.*
- HYDNUM fasciatum Peck, *syn. of Phellodon fasciatus q. v.*
- HYDNUM fennicum Sacc., *syn. of Sarcodon fennicus q. v.*
- HYDNUM ferrugineum Fr., *syn. of Hydnellum sanguinarium q. v.*
- HYDNUM flabelliforme Berk., *syn. of Steccherinum rhois q. v.*
- HYDNUM flavum Berkeley, *syn. of Grandinioides flavum q. v.*
- HYDNUM floriforme Schaeffer, *syn. of Hydnellum floriforme q. v.*
- HYDNUM fuligineo-violaceum Kalch., *syn. of Sarcodon fuligineo-violaceus q. v.*
- HYDNUM graveolens Delastre, *syn. of Phellodon graveolens q. v.*
- HYDNUM graveolens subzonatum Peck, *syn. of Phellodon coraceo-membranaceus q. v.*
- HYDNUM hybridum Pers., *syn. of Hydnellum floriforme q. v.*
- HYDNUM imbricatum L., *syn. of Sarcodon imbricatus q. v.*
- HYDNUM Key to species. [Banker.] Mem. Torr. Bot. Club, 12:106. 13 June 1906.
- HYDNUM laciniatum Leers, *syn. of Hericium laciniatum q. v.*
- HYDNUM niger Fries, *syn. of Phellodon niger q. v.*
- HYDNUM ochraceum Pers., *syn. of Steccherinum ochraceum q. v.*
- HYDNUM parasiticum Pers., *syn. of Steccherinum strigosum q. v.*
- HYDNUM plumarium B. & C. in Grev. non Jour. Linn. Soc. *syn. of Steccherinum ochraceum q. v.*
- HYDNUM plumarium B. & C. Jour. Linn. Soc. 10:324 [non H. plumarium B. & C. Grev. 1:97], *syn. of Steccherinum plumarium q. v.*

- HYDNUM pulcherrimum B. & C., *syn. of Steccherinum pulcherrimum q. v.*
- HYDNUM putidum Atkinson, *syn. of Phellodon putidus q. v.*
- HYDNUM ramosum Bull., *syn. of Hericium laciniatum q. v.*
- HYDNUM rhois Schw., *syn. of Steccherinum rhois q. v.*
- HYDNUM scabripes Peck, *syn. of Sarcodon scabripes q. v.*
- HYDNUM schiedermayeri Heuf., *syn. of Hericium croceum q. v.*
- HYDNUM scrobiculatum Fr., *syn. of Hydnellum scrobiculatum q. v.*
- HYDNUM septentrionale Fr., *syn. of Steccherinum septentrionale q. v.*
- HYDNUM spongiosipes Peck, *syn. of Hydnellum velutinum q. v.*
- HYDNUM stratosum Berk., *syn. of Leadia stratosa q. v.*
- HYDNUM strigosum Swartz, *syn. of Steccherinum strigosum q. v.*
- HYDNUM suaveolens Scop., *syn. of Hydnellum suavelens q. v.*
- HYDNUM suberosum aurantiacum Batsch, *syn. of Hydnellum floriforme q. v.*
- HYDNUM tinctorium Lloyd, *syn. of Echinodontium tinctorium q. v.*
- HYDNUM tomentosum L., *syn. of Phellodon tomentosus q. v.*
- HYDNUM vellereum Peck, *syn. of Phellodon vellereus q. v.*
- HYDNUM velutinum Fr., *syn. of Hydnellum velutinum q. v.*
- HYDNUM zonatum Batsch, *syn. of Hydnellum zonatum q. v.*
- HYMENOMYCETES, Culture Studies on Polymorphism of. George Richard Lyman. Proc. Boston Soc. Nat. Hist. 33:125-209. Pl. 18-26. Feb. 1907.
- HYMENOPSIS hydrophila Sacc. n. sp. in foliis emortuis Typhae latifoliae. Jour. Mycol. 13:47. Mar. 1907.
- HYPODERMA smilacis (Schw.) Sacc. *syn. of Gloniopsis smilacis q. v.*
- HYSTERIUM smilacis Schw., *syn. of Gloniopsis smilacis q. v.*
- HYSTEROGRAPHIUM smilacis (Schw.) E. & E. *syn. of Gloniopsis smilacis q. v.*
- INDEX to North American Mycology. W. A. Kellerman. Jour. Mycol. 13:85-7. Mar. 1907.
- KAUFFMAN, C. H. Unreported Michigan Fungi, from Petoskey, Detroit and Ann Arbor, for 1905. Mich. Acad. Sci. Meeting 1906, 8:26-37. Date of issue?



- KELLERMAN, Karl F., and Fawcett, Edna H. Movements of certain Bacteria in soils. [Abstract.] *Science N. S.* 25:806. 24 May 1907.
- KELLERMAN, W. A. [Ed.] *Mycological Bulletin*, Vol. IV, Nos. 49-72, pp. 193-286. 1906.
- KELLERMAN, W. A. Notes from *Mycological Literature*. XXII. *Jour. Mycol.* 13:72-84. Mar. 1907.
- KEY to species of *Hericium*. [Banker.] *Mem. Torr. Bot. Club*, 12:113. 13 June 1906.
- KEY to Genera of *Hydnaceae*. [Banker.] *Mem. Torr. Bot. Club*, 12:103. 13 June 1906.
- KEY to species of *Hydnellum*. [Banker.] *Mem. Torr. Bot. Club*, 12:150-1. 13 June 1906.
- KEY to species of *Hydnum*. [6 species; Banker.] *Mem. Torr. Bot. Club*, 12:106. 13 June 1906.
- KEY to species of *Leaia*. [Banker.] *Mem. Torr. Bot. Club*, 12:175. 13 June 1906.
- KEY to the species of *Melanosporae*. [Morgan.] *Jour. Mycol.* 13:53. Mar. 1907.
- KEY to the species of *Phellodon*. [Banker.] *Mem. Torr. Bot. Club*, 12:165-6. 13 June 1906.
- KEY to species of *Sarcodon*. [Banker.] *Mem. Torr. Bot. Club*, 12:138-9. 13 June 1906.
- KEY to species of *Steccherinum*. [Banker.] *Mem. Torr. Bot. Club*, 12:124-5. 13 June 1907.
- LACTARIA Persoon [antedates *Lactarius* Fries nearly half century. *Burlingham*]. *Torreya*, 7:118. June 1907.
- LACTARIAE, Suggestions for the study of. Gertrude Simmons *Burlingham*. *Torreya*, 7:118-123. June 1907.
- LACTARIUS Fries, *syn. of Lactaria Persoon q. v.*
- LEAIA Banker, n. gen. *Hydnaceae*. *Mem. Torr. Bot. Club*, 12:175. 13 June 1906.
- LEAIA, Key to species. [Banker.] *Mem. Torr. Bot. Club*, 12:175. 13 June 1906.
- LEAIA *piperata* Banker n. sp. on very rotten stump in damp woods. *Mem. Torr. Bot. Club*, 12:175. 13 June 1906.
- LEAIA *stratosa* (Berk.) Banker n. n. [*Hydnum stratosum* Berkeley.] *Mem. Torr. Bot. Club*, 12:177. 13 June 1906.
- LECANACTIS *zahlbruckneri* Herre n. sp. rare on maritime rocks 50 to 75 feet above the sea, San Francisco. *Bot. Gaz.* 43:270. April 1907.

- LENTINUS pulcherrimus, A new Lentinus from Pennsylvania. David R. Sumstine. *Torreyia*, 7:60-61. March 1907.
- LENTINUS pulcherrimus Sumstine n. sp., growing on buried stick. *Torreyia*, 7:60. March 1907.
- LENTODIUM [between the Agaricaceae and the Polyporaceae, with no near relatives. Lyman]. *Proc. Boston Soc. Nat. Hist.* 33:192. Feb. 1907.
- LEPTODON auriscalpium Quélet, *syn. of Auriscalpium auriscalpium* q. v.
- LEPTODON ochraceum Quél., *syn. of Steccherinum ochraceum* q. v.
- LICHEN Distribution in the Santa Cruz Peninsula California. Albert W. C. T. Herre. *Bot. Gaz.* 43:267-273. April 1907.
- LICHENS, Some Additions to the Flora of Middlesex Co., Mass. [24 specimens of lichens.] Reginald Heber Howe, Jr. *Bryologist* 9:81-2. Sept. 1906.
- LICHENS of the Adirondack League Club Tract [list of 60 species]. Carolyn W. Harris. *Bryologist*, 10:64-6. July 1907.
- LICHEN Notes No. 4. A Study of Umbilicaria vellea and Umbilicaria spadochroa. G. K. Merrill. *Bryologist*, 9:83-7. Sept. 1906.
- LICHENS, Some, of Mt. Watatic, Massachusetts. [List of Species.] Reginald Heber Howe, Jr. *Bryologist*, 9:46-48. May 1906.
- LIQUIDAMBER styraciflua L., lumber, host to Ceratostomella capillifera Hedgcock n. sp. *An. Rep. Mo. Bot. Gar.* 1906, 17:71. 27 Sept. 1906.
- LIST of New York Fungi. [Onondago County.] F. L. Stevens. *Jour. Mycol.* 13:67-72. Mar. 1907.
- LITHOPHRAGMA parviflora (Hook.) Nutt. host to Puccinia lithophragma Holway n. sp. Holway's N. A. *Ured.* 1:51. 15 May 1906.
- LLOYD, C. G. *Mycological Notes*. No. 24, Dec. 1906. pp. 293-308. pl. 91-93. [Concerning Phalloids, &c.]
- LLOYD, C. G., Letter No. 10. Paris, July, 1906, pp. 1-8. Pamphlet. [General remarks on Polyporii and comments on specimens sent by American correspondents.]
- LYCOPERDON septimum Lloyd n. sp. [specimens collected at Quito, Ecuador]. Lloyd's *Myc. Notes*, No. 24, p. 306. Dec. 1906.

- LYMAN, George Richard. Culture Studies on Polymorphism of Hymenomycetes. Proc. Boston Soc. Nat. Hist. 33:125-209. pl. 18-26. Feb. 1907.
- MCALPINE'S Studies of Australian Rusts [review.] J. C. Arthur. Jour. Mycol. 13:41-2. Mar. 1907.
- MACROPLODIA americana, see *Sphaeropsis americana*.
- MANINA cordiformis Scop., syn. of *Hericium erinaceus* q. v.
- MEDUSINA coralloides Chev., syn. of *Hericium coralloides* q. v.
- MEDUSINA coralloides Chev., syn. of *Hericium laciniatum* q. v.
- MEDUSINA patua Chev., syn. of *Hericium erinaceus*.
- MELANOSPORA parasitica. [Notes and Synonymy.] Guy West Wilson. Torreyea, 7:57-9. March 1907.
- MELONOSPORA parasitica L. Tul. & C. Tul. [includes *Sphaeroneema parasitica* L. Tul. Ann. Sci. Nat. and *Ceratostoma biparasiticum* E. & E.] Torreyea, 7:59. March 1907.
- MELANOSPORAE [Monograph and Key; Morgan. Jour. Mycol. 13:53. Mar. 1907.
- MELOTHRIA pendula L., host to *Puccinia meliothriae* Stevens n. sp. Bot. Gaz. 43:283. April 1907.
- MERRILL, G. K. Lichen Notes No. 4. A Study of *Umbilicaria vellea* and *Umbilicaria spadachroa*. Bryologist, 9:83-7. Sept. 1906.
- MICHENERA artocreas B. & C. [the imperfect form of *Corticium subgiganteum* Berk. Lyman]. Proc. Boston Soc. Nat. Hist. 33:151. Feb. 1907.
- MICHIGAN Fungi from Petoskey, Detroit, and Ann Arbor for 1905. Unreported. C. H. Kauffman. Mich. Acad. Sci. Meeting 1906, 8:26-37. Date?
- MICROCHLOA indica (L.) Kuntze, host to *Ustilago microchloae* Griffiths n. sp. Bull. Torr. Bot. Club, 34:207. April 1907.
- MOORE, C. L. Myxomycetes of Pictou County, N. S. [33 species.] Pictou Academy Bull. 1:11-16. June 1906.
- MORGAN, A. P. North American species of Agaricaceae [Monograph], the Melanosporae [Synopsis of Genera *Psathyrella* 20 species, and *Panaeolus* 15 species]. Jour. Mycol. 13:53-62. Mar. 1907.
- MT. Watatic Lichens, Massachusetts, Some. [List of Species.] Reginald Heber Howe, Jr. Bryologist, 9:46-48. May 1906.
- MOVEMENTS of Certain Bacteria in the Soil. Karl F. Kellerman and Edna H. Fawcett. [Abstract.] Science N. S. 25:806. 24 May 1907.



- MUHLENBERGIA schaffneri elongata, host to Tilletia muhlenbergiae Clinton n. sp. [Mexico.] N. A. Flora, 7:49. 4 Oct. 1906.
- MYCOLOGICAL Bulletin [W. A. Kellerman, E.] Vol. IV, Nos. 49-72, pp. 193-286. 1906.
- MYCOLOGICAL Notes. C. G. Lloyd. No. 24. Dec. 1906, pp. 293-308. Pl. 91-39. [Concerning Phalloids, etc.]
- MYXOMYCETES of Pictou County, N. S. [33 species.] C. L. Moore. Pictou Academy Bull. 1:11-6. June 1906.
- NECTRIA cinnabarina (Tode) Fries, *syn. of Nectria purpurea* q. v.
- NECTRIA purpurea (Linn.) Wilson & Seaver n. n. [Tremella purpurea Linn.; Tubercularia vulgaris Tode; Sphaeria cinnabarina Tode; Nectria cinnabarina (Tode) Fries]. Jour. Mycol. 13:51. Mar. 1907.
- NEW Fungi from New York. P. A. Saccardo. Jour. Mycol. 13:45-8. Text figs. Mar. 1907.
- NEW Lentinus from Pennsylvania — L. pulcherrimus. David R. Sumstine. Torrey, 7:60-61. March 1907.
- NEW Species, see Griffiths, David, Concerning. . . .
- NEW YORK Fungi [Onondago County] F. L. Stevens, see *List of*.
- NITROPHILA occidentalis, host to Aecidium sarcobati Peck. [Griffiths.] Bull. Torr. Bot. Club, 34:210. April 1907.
- NORTH American Species of Agaricaceae [Monograph], the Melanosporae [Synopsis of Genera Psathyrella 20 species, and Panaeolus 15 species]. A. P. Morgan. Jour. Mycol. 13:53-62. Mar. 1907.
- NORTH American Uredineae. E. W. D. Holway. Vol. I. Part III, pp. 57-80. Pl. 24-34. 10 May 1907.
- NORTH American Uredineae. E. W. D. Holway. Part II, 1:53-56. Pl. 11-23. 15 May 1906.
- NOTAE Mycologicae [New Species, North American]. P. A. Saccardo. Ann. Mycolog., 5:176-9. April [issued 15 May] 1907.
- NOTES from Mycological Literature, XXII. W. A. Kellerman. Jour. Mycol., 13:72-84. Mar. 1907.
- ORTON, W. A., Plant Diseases in 1906. Yearbook U. S. Dept. Agr. 1906:409-508. 1907.
- PANAEOLUS [Monograph; Morgan], see *North American Species of Agaricaceae* [Monograph].

- PANAEOLUS carbonarius Batsch [Agaricus retirugis Fries; Morgan]. Jour. Mycol, 13:61. Mar. 1907.
- PANICUM caespitosum Swartz, host to Sorosporium ovarium Griffiths n. sp. [Mexico.] Bull. Torr. Bot. Club, 34:209. April 1907.
- PANICUM paspaloides, host to Ustilago rickerii Clinton n. sp. [Cuba.] N. A. Flora, 7:11. 4 Oct. 1906.
- PANICUM rothboellioides, host to Sphacelotheca diplospora glabra Clinton n. var. [Cuba.] N. A. Flora, 7:27. 4 Oct. 1906
- PANICUM sp., host to Sphacelotheca diplospora verruculosa Clinton n. var. [Mexico.] N. A. Flora, 7:27. 4 Oct. 1906.
- PENIOPHORA candida (Persoon) Lyman n. sp. [Aegerita candida Pers.], on pieces of pine boards, and fallen branches of Alnus, Populus, Acer, etc., in wet places; hymenium always accompanied by the conidial form. Proc. Boston Soc. Nat. Hist., 33:168. February 1907.
- PEZIZA flava Swartz, *syn. of Grandinioides flavum q. v.*
- PHAEODON atroviride Earle, *syn. of Sarcodon atroviride q. v.*
- PHAEODON aurantiacus Schroet., *syn. of Hydnellum floriforme q. v.*
- PHAEODON fennicus Hennings, *syn. of Sarcodon fennicus q. v.*
- PHAEODON ferrugineus Schroet., *syn. of Hydnellum sanguinarium q. v.*
- PHAEODON imbricatus Schroet., *syn. of Sarcodon imbricatus q. v.*
- PHAEODON zonatus Schroet. *syn. of Hydnellum zonatum q. v.*
- PHALLOIDS, concerning. Mycological Notes. C. G. Lloyd. No. 24, Dec. 1906. pp. 293-308. pl. 91-93.
- PHELLODON alboniger (Peck) Banker n. n. [Hydnum albonigrum Peck.] Mem. Torr. Bot. Club, 12:167. 13 June 1906.
- PHELLODON coriaceo-membranaceus (Schw.) Banker n. n. [Hydnum coriaceo-membranaceum Schw.; H. graveolens subzonatum Peck.] Mem. Torr. Bot. Club, 12:172. 13 June 1906.
- PHELLODON cyathiformis Karsten, *syn. of Phellodon tomentosus q. v.*
- PHELLODON delicatus (Schw.) Banker n. n. [Hydnum delicatum Schw. non Klotsch.] Mem. Torr. Bot. Club, 12:173. 13 June 1906.
- PHELLODON ellisianus Banker n. sp. on the ground in wood road. Mem. Torr. Bot. Club, 12:174. 13 June 1906.

- PHELLODON fasciatus (Peck) Banker n. n. [Hydnum fasciatum Peck.] Mem. Torr. Bot. Club, 12:171. 13 June 1906.
- PHELLODON graveolens (Delast.) Banker n. n. [Hydnum graveolens Delastre.] Mem. Torr. Bot. Club, 12:169. 13 June 1906.
- PHELLODON, Key to the species. [Banker.] Mem. Torr. Bot. Club, 12:165-6. 13 June 1906.
- PHELLODON niger (Fr.) Karst. [Hydnum niger Fries; Hydnelum Karsten; Calodon niger Quélet.] Mem. Torr. Bot. Club, 12:166. 13 June 1906.
- PHELLODON putidus (Atkin.) Banker n. n. [Hydnum putidum Atkinson.] Mem. Torr. Bot. Club, 12:170. 13 June 1906.
- PHELLODON tomentosus (L.) Banker n. n. [Hydnum tomentosum L.; Hydnum cyathiforme Schaeffer non Bulliard; Hydnelum cyathiforme Karsten; Phellodon cyathiformis Karsten; Calodon cyathiformis Quélet.] Mem. Torr. Bot. Club, 12:171. 13 June 1906.
- PHELLODON vellereus (Peck) Banker n. n. [Hydnum vellereum Peck.] Mem. Torr. Bot. Club, 12:168. 13 June 1906.
- PHYSARUM cinereum Pers., A Grass-killing Slime Mould. John W. Harshberger. Proc. Am. Philosophical Soc. Philadelphia, 45:271-3. Oct.-Dec. 1906.
- PHYTOPHTHORA infestans Mont. and Plasmopara cubensis (B. & C.) Humph. in Ohio, On the occurrence of. A. D. Selby. Ohio Nat. 7:79-85. Feb. 1907.
- PINUS echinata, sapwood of lumber, host to Ceratostomella Schrenkiana Hedgcock n. sp. An. Rep. Mo. Bot. Car. 1906, 17:67. 27 Sept. 1906.
- PLANT Diseases. Report of Assistant in Botany and Horticulture. H. S. Fawcett. Fla. Agr. Exp. Sta. Ann. Rep. 1906; XXI-XXVII. 1907.
- PLANT Diseases in 1906. W. A. Orton. Yearbook U. S. Department of Agriculture 1906: 409-508. 1907.
- PLANT-TUMOR of Bacterial Origin [B. tumefaciens n. sp.]. Erwin F. Smith and C. O. Townsend. Science N. S. 25:671-2. 26 April 1907.
- PLASMOPIARA cubensis (B. & C.) Humph. and Phytophthora infestans Mont. in Ohio, On the occurrence of. A. D. Selby. Ohio Nat. 7:79-85. Feb. 1907.
- PLEURODON auriscalpium Karsten, syn. of Auriscalpium auriscalpium q. v.



- PLEOSPHERIA fairmaniana Sacc. n. sp. in ligno carioso indurato Ulmi americanae. Jour. Mycol. 13:45. Mar. 1907.
- POLYGONUM newberryi, host to Ustilago punctata Clinton n. sp. N. A. Flora, 7:23. 4 Oct. 1906.
- POLYMORPHISM of Hymenomycetes, Culture Studies on. George Richard Lyman. Proc. Boston Soc. Nat. Hist 33:125-209. pl. 18-26. Feb. 1907.
- PRESCOT, C. S., Secretary Society of Bacteriologists. 8th Annual Meeting. [Titles and abstracts of papers]. Science N. S. 25:805-820. May 24, 1907.
- PRUNUS serotina, host to Zygodemus avellaneus Sacc. n. sp. Jour. Mycol. 13:48. Mar. 1907.
- PSATHYRELLA [Monograph; Morgan], see *North American Species of Agaricaceae* [Monograph]. . . .
- Puccinia arabicola E. & E. Jour. Mycol. 6:119, 191, [is Puccinia plumbaria Peck on Phlox not Arabis. Holway]. Holway's N. A. Uredin. 1:47. 15 May 1906.
- Puccinia euphorbiae intumescens Syd., syn. of Puccinia intumescens q. v.
- Puccinia eurotiae Griffiths n. sp. on Eurotia lanata Moq. Bull. Torr. Bot. Club, 34:210. April 1907.
- Puccinia glabella Holway n. sp. on Boissduvalia glabella (Nutt). Walp. Holway's N. A. Uredin. 1:76. 10 May 1907.
- Puccinia intumescens Holway, new comb. [Puccinia euphorbiae intumescens Syd.] Holway's N. A. Uredin. 1:60. 10 May 1907.
- Puccinia lithophragmae Holway n. sp. on Lithophragma parviflora (Hook.) Nutt. Holway's N. A. Ured. 1:51. 15 May 1906.
- Puccinia ludwigiae (E. & E.) Holway n. comb. [Aecidium ludwigiae E. & E.; Puccinia nesaeae E. & E.; Dicaeoma nesaeae Kuntze; Allodus nesaeae Artb.] Holway's N. A. Uredin. 1:72. 10 May 1907.
- Puccinia melothriae Stevens n. sp. on Melothria pendula L. Bot. Gaz. 43:283. April 1907.
- Puccinia on Melothria [P. melothriae n. sp.]. F. L. Stevens. Bot. Gaz. 43:282-3. April 1907.
- Puccinia nesaeae E. & E., syn. of Puccinia ludwigiae q. v.
- Puccinia obliqua B. C. Jour. Linn. Soc. 10:356, 1869, [appears to be P. lateritia B. & C. on some of the Rubiaceae. Holway]. Holway's N. A. Uredin. 1:43. 15 May 1906.

(To be continued.)

## NOTES FROM MYCOLOGICAL LITERATURE. XXV.

W. A. KELLERMAN.

**Moore, C. L.**

In Bulletin Vol. I, No. 1, Pictou Academy, N. S., there is given a general account of the Myxomycetes and a list of 33 "Myxomycetes of Pictou County."

**Magnus, P.**

An interesting article appeared in the Berichte der Deutschen Botanischen Gesellschaft, Jahrgang 1904, Band XXII, Heft 7, entitled "Puccinia Ruebsaumeni P. Magn. n. sp., eine einen einjaehrigen Hexenbesen bildende Art." The species has heretofore been confused with or included in Puccinia caulicola Schneid. (P. Schneideri Schroet.) from which it differs in its larger spores; it is restricted to Origanon vulgare. A plate of figures shows the hexenbesen, mycelium with haustoria *in situ* and the teleutospores.

**Fawcett, H. S.**

In Report in Botany and Horticulture (Florida Agricultural Station, An. Rep. 1906) Mr. Fawcett gives an account of the important diseases for the year — Colletotrichum lindemuthianum (Sacc. & Magn.) Bri. & Cav.; Alternaria brassicae var. nigrescens Pegl.; Cercospora apii Hals; Cladosporium citri Penz.; Colletotrichum gloeosporioides Penz.; Pseudoperonospora cubensis (B. & C.) Rost.; Gloeosporium mangiferae, &c.

**Saccardo, P. A.**

In "Notae Mycologicae," Series IX, Annales Mycologici for April, 1907, the author describes 7 new species, one of which is North American, namely, Tuberculina davisiana Sacc. et Trav. sp. n.; hab. in foliis adhuc vivis Salicis cordatae.

**Griffiths, David.**

Dr. Griffiths reports "Concerning some West American Fungi" in the Bulletin of the Torrey Botanical Club, April, 1907. They are species of his personal collecting for the past four or five years, new or worthy of record. The new species are: Sclerospora farlowii; Ustilago microchloae, Sorosporium ovarium, Urocystis sophiae, Aecidium cannonii, Puccinia eurotiae.

**Campbell, Douglas Houghton.**

Chapter VI in A University text-book of Botany, 2nd Edition, 1907, is devoted to the Fungi. The general character of the Subkingdom Fungi is given briefly; similarly the Structure and Affinities are discussed. The bulk of the chapter is concerned

with the classification — the author recognizing the usual three classes — the Phycomycetes including the subclasses Omycetes and Zygomycetes; the two classes constituting the Eu-mycetes; Ascomycetes composed of Hemiascineae, and Euasceae; and Basidiomycetes composed of the three sub-classes; Hemibasideae, Protobasidiomycetes, and Autobasidiomycetes. Five pages are devoted to the Lichens and one page to Bibliography. Numerous text figures are used.

### **Rolfs, F. M.**

Experiments in reciprocal inoculation with pure cultures shows that *Cytospora rubescens* Nitschke is the pycnidial stage of *Valsa leucostoma* — a fungus well known in Europe, Australia and the United States. "Professor F. C. Stewart of the New York Agricultural Experiment Station, was the first American to call attention to the parasitic nature of the fungus." At the Missouri State Fruit Experiment Station it is an active parasite attacking the twigs, limbs and trunk of the peach, plum, apricot and cherrytrees. This is reported under the head of "Dieback of the Peach Trees" in *Science* of July 19, 1907.

### **Lauterborn, Robert.**

In Heft 5, Band XXV, 26 June 1907, an account is given of "Eine neue Gattung der Schwefelbakterien (*Thioploca schmidlei* nov. gen. nov. spec.)" — belonging to the family Beggiotoaceae, occurring in Untersees des Bodensees in der Gegend von Ermatingen, in 15-20 m. Tiefe das Innere des kalkreichen Grundschlücks durchziehend.

### **Zeitschrift fuer Pflanzenkrankheiten, XIV. Band, 1904.**

The principal mycological articles of this volume are as follows: Ueber die Botrytis-Krankheit der Tulpen, H. Klebahn; Ueber *Trichothecium roseum* Link, als Ursache der Bitterfaule von Fruechten, K. S. Iwanoff; Der Rost des Getreides in Schlesien im Sommer 1903, W. Remer; Beitrage zur Kenntniss des Pilzes in den Wurzelanschwellungen von *Alnus incana*, C. G. Bjoerkenheim; Die *Peronospora-recte* *Pseudoperonospora* Krankheit der Melonen und Gurken in Ungarn, Prof. Linhart; Eine wichtige *Gloeosporium*-Krankheit der Linden, R. Lambert; Ueber den klimatisch-biologischen Zusammenhang einer Reihe Getreidekrankheiten waehrend der letzten Jahre, J. R. Jungner.

### **Zeitschrift fuer Pflanzenkrankheiten, XV. Band, 1905.**

We have to record the following titles: Pilzkrankheiten in Indien im Jahre 1903, E. J. Butler; Die schaedlichsten Insecten und Pflanzenkrankheiten welche an den Kultuzpflanzen in Bulgarien waehrend des Jahres 1903 geschadigt haben; Kulturversuche mit Rostpilzen, XII Bericht, 1903 und 1904, H. Klebahn;



*Helminthosporium gramineum* Rabenh. und *Pleospora trichostoma* Wint. Fritz Noack; Die Schlerotienkrankheit bei den Forsythien, A. Osterwalder.

### **Burlingham, Gertrude Simmons.**

In the February Bulletin of the Torrey Botanical Club (1907) Miss Burlingham gives an account of "Some *Lactarii* from Windham County, Vermont," based on collections made mostly near Newfane, elevation about 500 meters, from the middle of July to the middle of September. She describes the following: *Lactarius aspidioides* Burlingham n. sp.; *L. bensleyae* Burlingham n. sp.; *L. isabellinus* Burlingham n. sp.; *L. minusculus* Burlingham n. sp.; *L. nitidis* Burlingham n. sp.; and raises *Pecki* var. *oculatus* (of *L. subdulcis*) to specific rank; besides listing three dozen or more other species. The article concludes with an excellent key to the species of Vermont.

### **Peck, Charles Horton.**

Dr. Peck describes 20 new species in the Bulletin of the Torrey Botanical Club for February 1907. Over half of them are Agarics, the others of various groups. Of the specially interesting forms may be noted *Hydnum sulcatipes* with stem like some species of *Helvella*, and *Peckiella hymenioides* on *Lactarius uvidus*, externally similar to *P. hymenii*. *Lentinus ventricosus* Peck. Bull. Torr. Bot. Club, 23:414, 1896, is transferred to the genus *Armillaria*.

### **Wilson, Guy West.**

A monograph of the genus *Albugo* is published in the February No. of the Bulletin of the Torrey Botanical Club, under the title "Studies in North American Peronosporales — The genus *Albugo*." A brief general account of *Albugo* [*Albugo* (Pers.) S. F. Gray, Nat. Arr. Brit. Pl. 1:540. 1821 *Uredo* § *Albugo* Pers. Syn. Meth. Fung. 223. 1881; *Cystopus* Lév. Ann. Sci. Nat. III, 8:371. 1847] as represented in North America is given, followed by a key to the species. Synonymy in full, with citations, is printed; also a complete list of the hosts arranged alphabetically under families. One new species is described — *A. occidentalis*; and species excludendae are *Cystopus euphorbiae* Cke. & Mass. and *C. salsolae* and *C. schlechteri* P. Sydow. Oospores of nine of the thirteen species are illustrated by figures.

### **Shear, Cornelius Lott.**

For some time Mr. Shear has been studying the diseases of the Cranberry during which time he has encountered many new fungi and his interesting paper recently published in the Bulletin of the Torrey Botanical Club (June), entitled "New Species of Fungi", deals mainly with the vacciniicolous species. He

describes 20 to 30 new species and proposes three new genera, two of the latter belonging to Sphaeropsidales (*Plagiorhabdus*, and *Bothrodiscus*), and one to the Ascomycetae (*Acanthorhynchus*). The generic names are significant: the first from *plagios*, oblique, and *rhabdos*, rod in allusion to the oblique appendage of the spores; the second from *bothros*, pit, and *discos*, disk; and the third from *acanthos*, thorn, and *rhynchos*, beak, suggested by the spiny beak.

### Journal of Mycology, Vol. 13, July, 1907.

The table of contents of this No. is as follows: Sumstine, *Polyporus Pennsylvanicus* Sp. Nov; Sheldon, A Study of the Leaf-Tip Blight of *Dracaena Fragrans*; Durand, The Mycological Writings of Theodor Holmskjold and their relations to Persoon's *Commentatio*; Morgan, North American Species of Agaricaceae; Ricker, Third Supplement to New Genera of Fungi; Kellerman, Index to North American Mycology; Kellerman, Notes from Mycological Literature XXIV; Editor's Notes.

Unintentionally the Index to the New Genera of the Third Supplement was omitted when the last installment was printed. It is therefore inserted below.

### INDEX TO THIRD SUPPLEMENT TO NEW GENERA

- |                     |                      |                     |
|---------------------|----------------------|---------------------|
| Acerbiella, 64      | Eurychasma, 63       | Neottiopezis, 66    |
| Allodus, 119        | Eurychasmaceae, 63   | Nephlyctis, 122     |
| Ameris, 119         | Fairmania, 156       | Nyssopsora, 122     |
| Anaphysmene, 155    | Fioriella, 156       | Peronoeutypa, 66    |
| Argotelium, 119     | Gallacea, 154        | Peronoeutypella, 66 |
| Ascochytopsis, 155  | Gallowaya, 121       | Physopella, 123     |
| Bonanseja, 64       | Grandinioides, 154   | Platycarpium, 157   |
| Botryoconis, 155    | Hemispora, 156       | Polioma, 123        |
| Bubákia, 119        | Hirneolina, 154      | Polythelis, 123     |
| Capnodiella, 64     | Hysteridium, 156     | Pontisma, 63        |
| Ceratelium, 119     | Klebahnia, 121       | Prospodium, 123     |
| Chnoopsora, 120     | Lysospora, 121       | Pseudostegia, 157   |
| Cionothrix, 120     | Leaia, 154           | Pythiacystis, 64    |
| Colletomanginia, 65 | Leptomitella, 65     | Ramulaspera, 158    |
| Corynespora, 155    | Leptosphaerulina, 65 | Schoenhornia, 158   |
| Cryphonectria, 65   | Lindauopsis, 157     | Sirolpidium, 64     |
| Cystingophora, 120  | Listerella, 63       | Spirechina, 123     |
| Delastreopsis, 65   | Listerellaceae, 63   | Stictoclypeolum, 66 |
| Delitschiella, 65   | Macalpinia, 122      | Telospora, 124      |
| Dendroecia, 120     | Mapea, 122           | Tranzschelia, 124   |
| Dichaenopsis, 156   | Melampsoropsis, 122  | Trematovalsa, 66    |
| Dicheirinia, 121    | Melanobasidium, 157  | Trichofusarium, 158 |
| Discospora, 121     | Muchmoria, 157       | Trichophyma, 67     |
| Earlea, 121         | Mycorhynchus, 157    | Uromycopsis, 124    |
| Endothiella, 65     | Necium, 122          | Whetstonia, 154     |



# JOURNAL OF MYCOLOGY

*A Periodical Devoted to North American Mycology. Issued Bimonthly; January, March, May, July, September and November. Price, \$2.00 per Year. To Foreign Subscribers \$2.25. Edited and Published by*

*W. A. KELLERMAN, PH. D., COLUMBUS, OHIO.*

---

## EDITOR'S NOTES.

We have referred to the American Code of Botanical Nomenclature as recently published in the Bulletin of the Torrey Botanical Club, April, 1907, and made a suggestion in connection with the subject dealt with in Section I. As to Section II (formation of names), we have nothing to say except approbation, unless it be to note that Canon 8 does not require that the names for Subclasses and higher groups have the feminine form. All other group names, if we except the Genus, are required by preceding Canons to have the termination *-ae*. Moreover, such usage has found almost universal acceptance; we believe the 8th Canon could properly require this.

---

Professor Saccardo has taken the lead in many phases of classificatory and nomenclatural reform in Mycology. We desire to call attention to his "Disposizione e Nomenclatura" of mycological groups recently published in the *Bulletino della Societa Botanica Italiana*, which we will reproduce in full in the next Number of the JOURNAL. In this he is consistent as regards the matter of terminations referred to above; and he has made his groups conform to advanced nomenclatorial views. For example, the termination *-aceae* is used for family names, and *-ales* for the orders; all other terminations are in *-ae*. A brief excerpt in advance of the full list exhibits these points:

## REGNUM VEGETABILE: PLANTÆ.

Series CRYPTOGRAMAE (Lin. 1737) em.

Subseries: MYCETAE seu FUNGI (Juss. 1728) em.

Divisio: EUMYCETAE Eichler 1883 (Hyphomycetae Bref. 1887 non Mart.)

Subdiv. TELEOMYCETAE Sacc. 1902.

Classis I. BASIDIOMYCETAE (DeBy. 1862) em. (Basidioporeae Lév. 1837).

Subcl. I. EUBASIDIAE (Schroet. 1889) em.

Ordo I. HYMENIALES (Fr. 1821) em. nom., seu Hymenomycetae Fr.

Fam. I. AGARICACEAE Fr. 1825, &c.



# PRICE LIST.

---

Journal of Mycology, Vols. 8-12 (1902-1906) each.....	\$2 00
Journal of Mycology, Current Vol.....	2 00
Index to North American Mycology (authors, titles, subjects, new species and hosts, new names and synonyms) since Dec. 31, 1900. <i>Printed on one side of page.</i> Twelve parts have been issued. Price per part	25
Index to Uredineous Infection Experiments with List of Species and Hosts for North America, <i>printed on one side of page</i> .....	50
Portraits of Mycologists, reprinted from the Journal of Mycology (Peck, Ellis, Arthur, Burrill, Farlow, Tracy, Earle, Atkinson, Holway, Thaxter, Clinton, Magnus, Oudemans, Patouillard, Hennings, Rehm, Cooke, Saccardo, Dietel, Sydow, H., Sydow, P., Hollos, Bresadola, Massee, Salmon, Lindau, each.....	10
Ohio Fungi exsiccati. W. A. Kellerman. Ten fascicles, each .....	1 00
Mycological Bulletin. A popular leaflet of 8 pages, illustrated, 12 Nos. per year. Vol. II, 1904, Vol. III, 1905, Vol. IV, 1906, each 50 cents; bound in cloth, 75 cents. Current year .....	25
New Genera of Fungi. Published since 1900; the original descriptions reproduced with full citation. Annual Supplements are to follow. Printed on one side of page only. Price.....	75
First Supplement to New Genera of Fungi.....	50
Second Supplement to New Genera.....	50
Index to Journal of Mycology, volumes 1-10.....	75
Mycological Glossary .....	25
North American Species of Marasmius; a monograph by by A. P. Morgan.....	50
North American Species of Lepiota; by A. P. Morgan..	50
Cortinari; Key to Species; by C. H. Kauffman.....	25

Address: W. A. KELLERMAN, Columbus, Ohio.

